

Medication Use Instruction Sheet

使肺泰[®] 定量噴霧吸入器 Seretide[®] Evohaler

■ Name of medication content


Fluticasone 、 Salmeterol

■ Mechanism of action

Reduce airway inflammation and dilate airway, treat reversible respiratory tract obstruction.

■ Instructions for use

1. Remove cap on device opening.



2. Connect the broader end of the spacer to device opening (mouth piece)



3. Shake inhaler up down (until bottle base become cool), make sure the above mention parts are connected tightly.



4. Breathe out first before breathing in.



5. Hold the mouth piece between your lips, press the inhaler once, at the same time slowly and consistently breathe in deeply for at least 3-5 seconds. (Please try to press and breathe in at the same time to ensure medication efficacy)



6. Then hold your breath for 5-10 seconds, and slowly breathe out.



Hold breath

This sheet does not content complete drug information, if there is any enquiry, please contact your doctor or pharmacist.



以病家為尊、以同仁為重、以北醫為榮

■ Precautions

1. Please gargle your mouth after each use, because mouth gargling can reduce the occurrence of side effect of lowered hoarse voice and oral thrush.
2. If a second dose is required, please inhale the second dose at an interval of 1-2 minutes after completion of the first inhalation.

■ How to clean?

- Clean the inhaler at least once a week. Separate the inhaler (plastic outer cover) and the medication bottle (metal bottle), wash the inhaler with water, dry it right after washing, avoid placing at overheated place (like under the sun), no need to air dry with hairdryer. Do not place medication bottle into water.

References :

Drug Instruction Sheet 、 Micromedex

- Telephone : (02) 2737-2181 # 8130
- Producer / Date : Department of Pharmacy / 105.12.27
- PFS-7200-9006

This sheet does not content complete drug information, if there is any enquiry,
please contact your doctor or pharmacist.



以病家為尊、以同仁為重、以北醫為榮