Medication Use Instruction Sheet Ultibro®Breezhaler


Medication Name

Indacaterol 150 mcg, glycopyrronium 50mcg

Indication and Mechanism

Used for Chronic Obstructive Pulmonary Disease (COPD). Can relieve wheezing, coughing, and shortness of breath by opening up the airways.


Directions



- 1. Remove cap (Picture 1).
- Tilt the mouthpiece to open (Picture 2).
- 3. Remove capsule from blister and place in chamber (Picture 3-5).
- Close mouthpiece until it clicks (Picture 6).
- Press side piercing buttons in once and release (Picture 7-8). (Do not shake.)
- 6. Breathe out gently, away from inhaler (Picture 9).
- Close lips around mouthpiece to form good seal.
- Breathe in slowly and steadily. The capsule will vibrate as you breathe (Picture 10).
- 9. Breathe in as long as possible
- Hold breath for about 5 seconds, or as long as possible. Remove inhaler from mouth (Picture 11).

Handout is for reference only Please refer to your doctor for specific instructions.

以病家為尊、以同仁為重、以北醫為榮



- 11. Breathe out gently and away from inhaler.
- 12. Open mouthpiece and remove used capsule (Picture 13).
- 13. If there is still drug in the capsule, repeat all steps starting from step 3.
- 14. Clean the chamber of the breezhalor.
- 15. Close mouthpiece and cap.

Be careful of...

- Not piercing the capsule
- Piercing the capsule more than once
- Not using a new capsule for each dose
- Not breathing in forcefully enough

What if I miss a dose?

Immediately take a dose within the same day. Do not use more than once a day.

Cleaning Instructions

Clean the inhaler once a week. Wipe the mouthpiece with a clean alcohol pad or tissue.

References :

- 1. Micromedex
- 2. Manufacturer's label
- Contact Number : (02)2737-2181 ext. 8130 , 8132
- Department/Date : Pharmacy/107.10.19
- PFS-7200-9024

Handout is for reference only Please refer to your doctor for specific instructions.

以病家為尊、以同仁為重、以北醫為榮