Medication Use Instruction Sheet 帝舒满®定量噴霧吸入器 Duasma®HFA

Name of medication content


Budesonide 200 mcg/Puff

Mechanism of action

Reduce airway inflammation, and can effectively reduce the over sensitive condition of the airway.

Instructions for use

- 1. Remove the cover of the device.
- 2. Connect the broader end of the spacer to the mouth piece of the device.
- 3. Shake inhaler up down (until the bottom of the bottle becomes cool), at the same time, ensure the above mentioned parts are tightly connected.
- 4. Breathe out first before breathing in.
- 5. Hold the mouth piece in between your lips, press the inhaler once, at the same time slowly and consistently breathe in deeply for at least 3-5 seconds. (Please try to press and breathe in at the same time to ensure medication efficacy.)
- 6. Hold your breath for 5-10 seconds, and then slowly breathe out.



This sheet does not content complete drug information, if there is any enquiry, please contact your doctor or pharmacist.











Precautions

- 1. Please gargle your mouth after each use, because mouth gargling can effectively reduce the occurrence of lowered horsy voice and oral thrush side effects.
- 2. If a second dose is to be inhaled, please inhale the second dose at an interval of 1-2 minutes after the completion of first inhalation action.

How to clean?

Clean the inhaler at least once a week. Separate the inhaler (plastic outer cover) and the medication bottle (metal bottle), the inhaler can be wash with water, dry it after washing, avoid placing it at area with overheated temperature (under the sun), and do not air dry with hairdryer. <u>Metal bottle cannot be placed into water</u>.

References: Drug Instruction Sheet 、 Micromedex

- Telephone : (02) 2737-2181 # 8130
- Producer / Date: Department of Pharmacy / 105.12, 111.10 修
- PFS-7200-9002_EN

This sheet does not content complete drug information, if there is any enquiry, please contact your doctor or pharmacist.

