Medication Use Instruction Sheet 吸必擴®氣化噴霧劑 Symbicort®Rapihaler

■ Name of medication content

Budensonide 160 mcg, Formoterol 4.5 mcg

■ Mechanism of action

Reduce airway inflammation and dilate airway, treat reversible respiratory tract obstruction.

■ How to use your medicine?









- Remove the cover of the device by 1. pressing the both side and shake it up and down gently for 5 seconds.
- 2. Press the canister several times until a mist coming out from the device. This is to ensure the device is ready to be used. You should do it every time when you use a new device.
- 3. Slowly breath out completely away from inhaler. Put mouthpiece between teeth without biting and close lips to form a good seal.
- Start to breathe in slowly through mouth 4. and, at the same time, press down firmly on canister. Continue to breathe in slowly and deeply.
- Hold breath for about 5 seconds or as long 5. as comfortable.
- 6. Breathe out gently away from inhaler.
- 7. If more than one dose is needed, after 30 seconds, repeat all steps starting from step 3-6.
- 8. Rinse your mouth with water 2 times and spit after inhaling the last dose to reduce the risk of side-effects.

This sheet is for your information; please follow your doctor's directions.




■ Precaution

- 1.Please rinse your mouth every time after use to avoid hoarse voice and local infection.
 - 2.Please space out for 30 seconds between two doses.
 - 3. Pleae follow the instruction strictly.
- 4.Please store in room temperature (below 50 $^{\circ}$ C) once remove the device out of the package. Avoid moisture and direct sunlight. (DO NOT FREEZE and DO NOT REFRIGERATE)

■ How to know when the medicine is running out?

When the arrow pointed at the yellow zone, it means you have 20 doses remaining. It is time to schedule your next doctor visit. When the arrow pointed at the red zone, there is no medicine left.



■ How to clean?

Please clean the mouth piece at least once a week. Please wipe it with alcohol pad or tissue; do not dry it with hairdryer. DO NOT PUT THE CANISTER IN TO WATER.

Reference:

- 1. Micromedex
- 2. Drug information
- Consultation Number: (02)2737-2181 EXT 8130, 8132
- Publisher /Date: Pharmacy /106.06, 111.10 修
- PFS-7200-9007 EN



This sheet is for your information; please follow your doctor's directions.

