


Medication Use Instruction Sheet

Instructions for Use of Eye Drops

■ Instructions for use

1. Clean both hands thoroughly with soap and water.
2. Remove contact lenses before applying medicated eye drops.
3. Gently press or pull down the lower eye lid of the eye directed by the physician for medication use to expose an opening.
4. Tilt your head backward, place the dropper tube or container at a 2-3 cm distance above the lower eye lid, look up, and drop the solution droplet.
5. Try not to get the dropper tube or container in contact with your finger, face or eyes.
6. Keep your eyes closed for at least 30 seconds to prevent medication from flowing out of the eye.
7. Use your finger to assert pressure on the inner corner of the eye for 1-2 minutes to prevent medication from entering through the nose.



Dropper not
in contact
with finger,
face, or eyes.

Gently pull
down lower
eye lid.

This sheet does not content complete drug information, if there is any enquiry,
please contact your doctor or pharmacist.



以病家為尊、以同仁為重、以北醫為榮

■ Precautions

1. Do not allow dropper tube or container opening to contact with any object.
2. Do not flush the dropper tube or container opening with water.
3. Eye drops suspension should be shaken to homogenous before use.
4. Normally an eye could accommodate one drop of solution, if a few types of medicated eye drops are to be used concomitantly, have 5-10 minutes interval between each eye drop use to allow adequate absorption of medication, then followed by the second type of medication.
5. If eye drops and eye ointment were to be use concomitantly, please use eye drops first, then use eye ointment 5 minutes afterward.
6. Do not use eye drops with crystal formation or color change.
7. Do not use left-over of used steroid eye drops (fungi grow easily).
8. Do not share the same bottle of eye drops with other individual.
9. Do not allow direct sunlight exposure.
10. Some eye drops solution may cause blurred vision, do not drive or operate dangerous machine before having clear vision.
11. If the disease condition worsens or if there is itchy or burning sensation lasting for a few minutes after using eye drops, seek treatment immediately.
12. Unless instructed by pharmacist or physician, do not store eye drops in the refrigerator.
13. Discard eye drops one month after opening.

■ Telephone : (02) 2737-2181 # 8130

■ Producer / Date : Department of Pharmacy / 105.12, 111.10 修

■ PFS-7200-9014_EN

This sheet does not content complete drug information, if there is any enquiry,
please contact your doctor or pharmacist.



以病家為尊、以同仁為重、以北醫為榮