

# Instruction for Use

## Trulicity® 1.5 mg/0.5 mL/pen

### ■ What is in Trulicity® and what does Trulicity® do?



Trulicity® pen contains an injectable medicine called dulaglutide, that can be used along with diet and exercise to lower blood sugar in patients with type 2 diabetes mellitus. Trulicity® is a **disposable, prefilled medicine** delivery device that is given **once a week** at any time during this day, with or without meals.

### ■ What does Trulicity® look like?

Dulaglutide is a colorless, clear, aqueous solution.

### ■ How to use Trulicity®

1. Take out the pen from the refrigerator. Check the medicine to make sure it is not cloudy, discolored or has particles in it.
2. Make sure your pen is **locked**. Pull the base cap straight off and throw it away in your household trash.
3. Clean the injection site with an alcohol swab. Place the clear base flat and firmly against your skin at the injection site.
4. **Unlock** by turning the lock ring; press and hold the green injection button and you will hear a loud click.
5. **Continue holding** the clear base firmly against your skin until you hear a second click. This happens when the needle starts retracting in about 5-10 seconds.
6. You will know your injection is complete when the gray plunger is visible, then remove the pen from your skin.
7. Put your used pens in a puncture-resistant container right away after use. Do **NOT** throw away pens in your household trash.



This sheet is for your information; please follow your doctor's directions



以病家為尊、以同仁為重、以北醫為榮

## ■ Precaution

1. **Rotate your injection sites** within the area you choose each week. You may use the same area of your body, but be sure to choose a different injection site in that area.
2. Do not mix insulin and Trulicity® together in the same injection.
3. If you miss a dose of Trulicity®, take the missed dose as soon as possible if there are at least 3 days (72 hours) until your next scheduled dose. If there are less than 3 days remaining, skip the missed dose and take your next dose on the regularly scheduled day. Do not take 2 doses of Trulicity® within 3 days of each other.
4. If you wish to switch your injection day, **there must be 3 days (72 hours) between doses.** If it has been fewer than 3 days since your last dose, wait until the following week to take your dose on your new injection day.

## ■ How to store your Trulicity® pen

1. Keep new pens in the refrigerator between 2 °C and 8 °C (36 °F and 46 °F). **Do not freeze.**
2. You may store your pen at room temperature below 30 °C (86 °F) for up to a total of 14 days.
3. Storage of your pen in the original carton is recommended. The pen has glass parts; handle it carefully.
4. Keep your Trulicity® pen out of the reach of children.

## ■ Possible side effects

1. Some most common side effects of Trulicity® are nausea, diarrhea, vomiting, decreased appetite, stomach (abdominal) pain.
2. Other possible and severe side effects may include inflammation of your pancreas (pancreatitis), low blood sugar (hypoglycemia), change in vision, allergic reactions, and thyroid tumors. Stop using Trulicity® and contact your healthcare provider about any symptom that bothers you or does not go away.

### References:

Package insert 、 Micromedex

- Contact number : (02) 2737-2181 EXT. 8130
- Last updated by the Department of Pharmacy on 110.01, 111.10 修
- PFS-7200-9035\_EN

This sheet is for your information; please follow your doctor's directions

