




---

# 食道癌診療指引

**Esophageal and  
Esophagogastric Junction  
Cancer**

---

# 食道團隊成員



# Pretreatment Workup


---

## WORKUP

- H &P
- Upper GI endoscopy and biopsy
- ChestCT with oral and IV contrast( if no contraindications )
- PET-CT evaluation
- CBC and Chemistry profile
- Endoscopic ultrasound(EUS)(M1 病人可免做 )
- Endoscopic mucosal resection is essential for the accurate staging of early stage cancers( T1a or T1b)
- Biopsy of metastatic disease as clinically indicated
- MSI by PCR/MMR by IHC, and PD-L1 testing( preferred for documented/suspected metastatic disease )
- HER2-neu testing if metastatic adenocarcinoma is documented/ suspected
- Next-generation sequencing (NGS), may be considered
- Bronchoscopy, if tumor is at or above the carina with no evidence of M1 disease
- Assign Siewert category
- Nutritional assessment and counseling
- Smoking cessation advice, counseling, and pharmacotherapy as indicated
- Screen for family history( preferred )

## Pretreatment Workup


### CLINICAL STAGE



\* Percutaneous gastrostomy tube (PCG) may be considered for patient with cervical esophageal tumors receiving definitive chemoradiation or for patients with marginally resectable disease.


## 《 Esophageal cancer guideline-2》

### Squamous cell carcinoma: primary treatment options for medically fit surgery patients




\* ER: endoscopic resection

## Squamous cell carcinoma: response assessment



## Squamous cell carcinoma: surgical outcomes


Patients **Have Not** Received Preoperative Chemoradiation



\*High risk factors: close circumferential/surgical margin, LVI, perineural invasion, extracapsular nodal extension

## Squamous cell carcinoma: surgical outcomes


Patients **Have** Received Preoperative Chemoradiation






《 Esophageal cancer guideline-6》

## Squamous cell carcinoma & adenocarcinoma : Non- surgical candidate




\* Poor prognostic features: positive margin(s), max. tumor diameter > 2cm, G2/3, LVI or more


## Squamous cell carcinoma & adenocarcinoma follow up – recurrence – palliative management



## Squamous cell carcinoma & adenocarcinoma : unresectable locally advanced, locally recurrent, or metastatic disease




## Adenocarcinoma: primary treatment for medically fit surgery patients



# 《 Esophageal cancer guideline-10》


## Adenocarcinoma: response assessment



## Adenocarcinoma: surgical outcomes

Patients Have Not Received  
Preoperative Chemoradiation  
or chemotherapy


Postoperative management



## 《 Esophageal cancer guideline-12》

### Adenocarcinoma: surgical outcomes

Patients Have Received  
Preoperative Chemoradiation  
or chemotherapy



## Principles of pathological review

- Pathology review 的目的包括：
  - Classification of tumor
  - Determine the extent of invasion
  - Establish status of cancer involvement of surgical margins
- 所有手術病理報告都應該依照食道癌 WHO 分類
- 所有手術病理報告都應該依 AJCC/UICC TNM 8<sup>th</sup> edition 分期
- 手術病理報告應包括下列項目
  - Histologic type
  - Histologic grade (G1: well differentiated; G2: moderately differentiated; G3: poorly differentiated)
  - Microscopic tumor extension
  - Margin status



## Principles of pathological review

- 病理報告根據標本的不同應包括下列項目：
- Biopsy: invasion, if present; high-grade dysplasia in Barrett esophagus; histologic type; Grade; Presence or absence of Barrett esophagus
- Endoscopic resection (ER): include all elements as for biopsy specimen plus the depth of tumor invasion; lymphovascular invasion (LVI), and the status of mucosal and deep margins
- Esophagectomy, without prior chemoradiation: include all elements as for ER specimen plus the location of the tumor midpoint in relation to the EGJ, whether the tumor crosses EGJ, lymph node status, and the number of lymph nodes recovered
- Esophagectomy, with prior chemoradiation :
  - the tumor sites should be thoroughly sampled, with submission of entire EGJ or ulcer/tumor bed for specimens without grossly obvious residual tumor
  - For pathology report, include all elements as for esophagectomy without prior chemoradiation, plus assessment of the treatment effect
  - Assessment treatment effect: 術前治療反應與預後相關。目前採用 The modified Ryan scheme in the CAP Cancer Protocol for Esophageal Carcinoma

Description	Tumor Regression Score
No viable cancer cells (complete response)	0
Single cells or rare small groups of cancer cells (near complete response)	1
Residual cancer with evident tumor regression, but more than single cells or rare small groups of cancer cells (partial response)	2
Extensive residual cancer with no evident tumor regression (poor or no response)	3

- HER2 狀態，微衛星不穩定性（MSI）狀態的分子測試，程序性死亡配體 1（PD-L1）表達和 NTRK 基因融合檢測被用於預測局部晚期，不可切除或轉移性食道癌和 EGJ 癌的臨床治療用藥選擇
- Assessment of overexpression or amplification of Her2 in Esophageal and EGJ Cancer
  - For patients with inoperable locally advanced, recurrent, or metastatic adenocarcinoma of esophagus or EGJ for whom trastuzumab therapy is being considered
  - Immunohistochemical criteria for scoring HER2/neu expression

	Surgical Specimen Expression Pattern, Immunohistochemistry	Biopsy Specimen Expression Pattern, Immunohistochemistry	HER2 Overexpression Assessment
0	No reactivity or membranous reactivity in <10% of cancer cells	No reactivity or no membranous reactivity in any cancer cell	Negative
1+	Faint or barely perceptible membranous reactivity in $\geq 10\%$ of cancer cells; cells are reactive only in part of their membrane	Cluster of five or more cancer cells with a faint or barely perceptible membranous reactivity irrespective of percentage of cancer cells positive	Negative
2+	Weak to moderate complete, basolateral or lateral membranous reactivity in $\geq 10\%$ of cancer cells	Cluster of five or more cancer cells with a weak to moderate complete, basolateral, or lateral membranous reactivity irrespective of percentage of cancer cells positive	Equivocal
3+	Strong complete, basolateral, or lateral membranous reactivity in $\geq 10\%$ of cancer cells	Cluster of five or more cancer cells with a strong complete, basolateral, or lateral membranous reactivity irrespective of percentage of cancer cells positive	Positive

- HER2 IHC is performed first, followed by FISH methods in cases showing 2+ (equivocal) expressions by IHC. Cases with HER2: CEP17 ratio  $\geq 2$  or an average HER2 copy number  $\geq 6.0$  signals/cell are considered positive by FISH.

## Principles of biomarker testing

- Mismatch Repair (MMR) Testing:
  - MMR by IHC should be considered on locally advanced, recurrent, or metastatic esophageal and EGJ cancer in patents who are candidate for treatment with PD-1 inhibitors
  - MMR status is assessed by IHC staining performed on formalin-fixed , paraffin-embedded (FFPE) tissue to measure levels of proteins involved in DNA mismatch repair including MLH1, MSH2, MSH6, and PMS2
  - MMR IHC interpretation
    - No loss of nuclear expression of MMR proteins: No evidence of deficient mismatch repair (low probability of MSI-H)
    - Loss of nuclear expression of one or more MMR proteins: deficient mismatch repair
- PD-L1 Testing:
  - PD-L1 IHC testing may be considered on locally advanced, recurrent, or metastatic esophageal and EGJ cancer in patents who are candidate for treatment with PD-1 inhibitors
  - Assessment of PD-L1 Protein Expression
    - Pembrolizumab as a second-line treatment option for esophageal SCC with PD-L1 expression levels by combined positive score (CPS) of  $\geq 10$ , and as a third-or subsequent-line treatment option for EGJ adenocarcinoma with PD-L1 expression levels by CPS  $\geq 1$ , as determined by an FDA-approved companion diagnosed test
    - CPS is determined by

$$\text{CPS} = \frac{\text{\# of PD-L1-positive cells (tumor cells, lymphocytes, macrophages)}}{\text{Total \# of tumor cells}} \times 100$$

## Principles of biomarker testing

- Assessment of *NTRK* gene fusions:
  - The FDA granted approval for the use of select TRK inhibitors for *NTRK* gene fusion-positive solid tumors
  - A two-step approach is used, which includes IHC first and confirmation of any positivity detected with IHC by Next generation sequencing (NGS)
  - TRK IHC as a screening tool:
    - IHC negative: No TRK expression
    - IHC positive: Detection of TRK expression, confirmation by NGS
  
- 若只有少量組織 (biopsy specimen) 可供檢測不同種類的 biomarker，建議臨床醫師開立檢測請在病理申請單註明，同一次病理組織切片除了 HE，應預留組織空白片，以避免因蠟塊多次處理，造成腫瘤細胞消耗

## Principles of Endoscopic staging and therapy

1. 食道鱗狀細胞癌病患，在初步確診時，必須由專精影像強化技術 IEE 的內視鏡醫師進行一次上消化道內視鏡精查，以確認是否同時存在其他源於黏膜的癌症及癌前病變。(work up)
2. 所有食道癌等病患，在進行手術或 CCRT 前，應會診胃腸科醫師討論是否施行內視鏡或外科性胃腸造口，以顧全病患治療前後的營養狀態。(principle of surgery)
3. 食道癌在接受治療前，除了已明確證實為 M1 的病人外（如 CT scan、PET scan），應進行 EUS 診斷。(work up)
4. 關於早期食道癌：適用內視鏡切除的絕對適應症為腫瘤深度侷限在 m1、m2 層，且 LN(-)；相對適應症為腫瘤深度侷限在 m3、sm1 層，且 LN(-)；無法辨明情況下，可考慮行 diagnostic ESD。
5. 如果在上述測試完成後有足夠的組織可用，可以考慮用 NGS 做第二次檢測。
6. 食道癌病患若 EUS 發現 LN(+)，可選擇性加做 FNA/FNB。
7. 食道癌 CCRT 之後，可選擇性加做 EUS 做 re-staging。(optional)
8. 姑息性的內視鏡治療（例如：金屬支架擴張術、氫氣電漿凝固術局部消融…等），應於食道癌多科團隊會議中提出討論後執行。

註一：執行上消化道內視鏡精查時，應備有擴大型內視鏡、魯格爾試液（Lugol's solution）、稀釋醋酸、影像強化光源（例如 NBI、BLI 等），由具早期癌判斷能力的內視鏡醫師執行並撰寫報告。

註二：巴瑞特氏食道病患，須例行接受上消化道內視鏡精查，以確認是否發生癌變，精查應由專精影像強化技術 IEE 的內視鏡醫師執行。（有 dysplasia 者至少每年一次，無 dysplasia 者至少每三年一次。）

註三：應由食道癌共識會議向三院相關癌症多科團隊提出診療建議「所有口腔癌（含舌癌）、頭頸癌、下咽癌、喉癌等病患，在初步確診時及術後至少每年，必須由專精影像強化技術 IEE 的內視鏡醫師進行一次上消化道內視鏡精查，以確認是否同時存有食道癌及癌前病變。」

## Principle of surgery

- 手術前之臨床分期應使用胸腹部電腦斷層，正子電腦斷層掃描 (PET-CT) 及內視鏡超音波，以評估可否切除
- 治療前所有病患皆需諮詢胸腔外科醫師，接受是否可以耐受食道切除之生理評估
- 所有生理狀態適合食道切除，且為局部可切除之胸腔 (超過環咽部 5 公分以下) 及腹腔食道癌患者，皆應考慮手術切除
- 食道交接處 (EGJ) 腺癌之病患皆應有 Siewert 分類
  - Siewert Type I: adenocarcinoma of the lower esophagus with the epicenter location located within 1cm to 5 cm above the anatomic EGJ
  - Siewert Type II: true carcinoma of the cardia with the tumor epicenter within 1 cm above and 2 cm below the EGJ
  - Siewert Type III: subcardial carcinoma with the tumor epicenter between 2 cm and 5 cm below the EGJ, which infiltrates the EGJ and lower esophagus from below
- Siewert type I 及 II 腫瘤應遵從食道癌及胃食道交接癌治療準則。Siewert type III 則視為胃癌，應遵從胃癌治療準則，然某些狀況下需增加部分食道切除，以獲取適當的之切除距離
- 腹腔內視鏡可以偵測放射線影像檢查無法找到之潛伏轉移病灶 (Occult metastatic disease) 尤其是 Siewert type II 及 III 腫瘤
- 腹膜積液細胞學檢查陽性與不良預後相關，應視為 M1 疾患。患者若為進展期之腫瘤，臨床分期 T3 或 N+ 疾患，應考慮接受腹腔內視鏡腹膜沖洗分期檢查 (laparoscopic staging with peritoneal washings)。
- 頸部食道癌或不超過環咽部 (cricopharygeal) 5 公分以上之食道癌應該以決定性放射化療 (definitive chemoradiation) 治療

## Principle of surgery

- 可切除的食道癌 Stage I-IVA(Locoregional disease, except T4b or unresectable N3) :
  - Tis or T1a 腫瘤 ( 可考慮 EMR+Ablation 或手術 )
  - T1b 腫瘤
  - T1-T3 腫瘤，即使局部淋巴腺已經轉移，多區域 (multi-station) 淋巴腺或巨大 (bulky) 淋巴腺轉移為手術切除之 relative contraindication，可否切除尚應考慮，如年齡、體能狀態、或治療反應等其他因素
  - T4a 腫瘤，侵犯至肋膜、心包膜或橫膈膜
- 不可切除的食道癌 Stage IVA(including T4b or unresectable N3) & IVB(metastatic disease)
  - cT4b 腫瘤：侵犯心臟、大血管、氣管、或鄰近器官包括肝、胰、肺及脾臟
  - 多區域 (multi-station) 淋巴腺或巨大 (bulky) 淋巴腺侵犯，大多數應視為不可切除，然而淋巴侵犯可否切除尚應考慮，如年齡、體能狀態、或治療反應等其他因素
  - EJG 腫瘤患者合併鎖骨上淋巴轉移應視為不可切除
  - 轉移第 IV 期之食道癌 ( 包括非區域淋巴腺 non-regional lymph node 轉移 )
- 食道切除方式之選擇取決於腫瘤位置、可選擇重建之器官，手術者之經驗及喜好，以及病患之喜好
- 病患在術前引導期間無法經口進食以維持營養者可考慮食道擴張或空腸造瘻 (J-tube)，優於胃造瘻術 ( 會妨礙往後胃重建手術時胃管之健全

## Principle of surgery

- 可接受之食道及胃交接腫瘤切除術式
  - Ivor Lewis esophagogastrectomy (開腹及右側開胸)
  - McKeown esophagogastrectomy (右側開胸、開腹及頸部吻合)
  - Minimally invasive Ivor Lewis esophagogastrectomy (腹腔鏡及微創右側開胸)
  - Minimally invasive McKeown esophagogastrectomy (胸腔鏡、微創開腹 / 腹腔鏡及頸部吻合)
  - Transhiatal esophagogastrectomy (開腹不經胸腔及頸部吻合)
  - Robotic minimally invasive esophagogastrectomy (機器手臂輔助微創術式)
  - Left transthoracic or thoracoabdominal approaches with anastomosis in chest or neck(左側開胸或胸腹聯合術式、胸腔或頸部吻合)
- 可接受之食道重建取代物
  - 胃 (優先選擇)
  - 大腸
  - 空腸
- 可接受之淋巴擴清方式
  - 標準方式
  - 擴展方式 (整體 En-bloc 方式)
- 至少需移除或評估 15 個以上之淋巴結以達到適當的淋巴分期, 若術前接受過放射化學治療, 雖適切需移除或評估之淋巴結數量仍然未知, 仍建議移除或評估 15 個以上之淋巴結
- 患者經過決定性放射及化學治療後, 仍有可切除之局部腫瘤, 且沒有遠端轉移, 可以考慮食道切除
- 食道切除, 內視鏡食道黏膜切除, 及其他燒灼術式, 應在高容量之食道治療中心由有經驗的外科醫師或內視鏡醫師執行



## Reference for principle of surgery

1. NCCN Practice Guidelines in Oncology, Esophageal Cancer, 2022 V4.
2. Hsu PK, Huang CS, Wang BY, Wu YC, Hsu WH. Survival Benefits of Postoperative Chemoradiation in Lymph Node-Positive Esophageal Squamous Cell Carcinoma. *Ann Thorac Surg* 2014;97:1734-41. (SCI)
3. Hsu PK, Huang CS, Wu YC, Chou TY, Hsu WH. Open versus Thoracoscopic Esophagectomy in Patients with Esophageal Squamous Cell Carcinoma. *World J Surg.* 2014;38:402-9. (SCI)
4. Hsu PK, Huang CS, Wang BY, Wu YC, Chou TY, Hsu WH. The Prognostic Value of the Number of Negative Lymph Nodes in Esophageal Cancer Patients after Transthoracic Resection. *Ann Thorac Surg* 2013;96:995-1001. (SCI)
5. Hsu PK, Chien LI, Huang CS, Hsieh CC, Wu YC, Hsu WH, Chou TY. Comparison of survival among neoadjuvant chemoradiation responders, non-responders and patients receiving primary resection for locally advanced oesophageal squamous cell carcinoma: does neoadjuvant chemoradiation benefit all? *Interact Cardiovasc Thorac Surg.* 2013;17:460-6. (SCI)
6. Wang BY, Liu CY, Lin CH, Hsu PK, Hsu WH, Wu YC, Cheng CY. Endoscopic Tumor Length Is an Independent Prognostic Factor in Esophageal Squamous Cell Carcinoma. *Ann Surg Oncol.* 2012;19:2149-58. (SCI)
7. Hsu PK, Wang BY, Huang CS, Wu YC, Hsu WH. Prognostic factors for post-recurrence survival in esophageal squamous cell carcinoma patients with recurrence after resection. *J Gastrointest Surg.* 2011;15:558-65. (SCI)
8. Hsu PK, Wang BY, Chou TY, Huang CS, Wu YC, Hsu WH. The total number of resected lymph node is not a prognostic factor for recurrence in esophageal squamous cell carcinoma patients undergone transthoracic esophagectomy. *J Surg Oncol.* 2011;103:416-20. (SCI)
9. Wang BY, Goan YG, Hsu PK, Hsu WH, Wu YC. Tumor length as a prognostic factor in esophageal squamous cell carcinoma. *Ann Thorac Surg.* 2011;91:887-93. (SCI)

## Follow Up

- History and Physical examination
  - Every 3 to 6 months for 1 to 2 years.
  - Every 6 to 12 months for 3 to 5 years, then annually.
- Clinical examination  
Optional as clinically indicated
  - Chemistry profile and CBC, as clinically indicated.
  - Imaging as clinically indicated.
  - Upper GI endoscopy and biopsy as clinically indicated.
  - Dilatation for anastomotic stenosis.
  - Nutritional assessment and counseling.

## 《 食道癌抗癌藥物治療指引 》

### Preoperative Chemoradiation Therapy of Esophageal Cancer Preferred Regimens

#### Paclitaxel + Carboplatin

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	50	1	QW	5	1
Carboplatin	AUC 2	1	QW	5	

#### Cisplatin + Fluoropyrimidine

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	75	1	Q4W	2(RT) + 2	2
5-FU	1000*	1-4	Q4W	2(RT) + 2	

\* Continuous infusion for 24 hours

藥品名 *	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	15	1-5	Q3W	2	5
5-FU	800*	1-5	Q3W	2	

\* Continuous infusion for 24 hours

藥品名 *	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	30	1	QW	5	44
Capecitabine	800 PO BID	1-5	QW	5	

### Oxaliplatin + Fluoropyrimidine

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	3(RT) + 3	2
Leucovorin	400	1	Q2W	3(RT) + 3	
5-FU	400	1	Q2W	3(RT) + 3	
5-FU	800*	1-2	Q2W	3(RT) + 3	

\* Continuous infusion for 24 hours

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1, 15, 29		For 3 doses	43
Capecitabine	625 PO BID	1-5	QW	5	

### Other Regimens

#### Cisplatin + Fluorouracil

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	100	1, 29	Q35D	1	4
5-Fu	1000	1-4, 29-32	Q35D	1	

**Irinotecan + Cisplatin**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Irinotecan	65	1, 8, 22, 29		1	6
Cisplatin	30	1, 8, 22, 29		1	

**Paclitaxel + Fluoropyrimidine**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	45	1	QW	5	7
5-FU	300*	1-5	QW	5	

\* Continuous infusion for 24 hours

**Perioperative Chemotherapy (Include EGJ)****Preferred Regimens****ECF modications**

藥品名 *	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Epirubicin	50	1	Q3W	6*	9
Cisplatin	60	1	Q3W	6*	
Capecitabine	625 PO BID	1-21	Q3W	6*	

\*3(pre-op) + 3(post-op)

藥品名 *	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Epirubicin	50	1	Q3W	6*	9
Oxaliplatin	130	1	Q3W	6*	
Capecitabine	625 PO BID	1-21	Q3W	6*	

\*3(pre-op) + 3(post-op)

### Oxaliplatin + Fluoropyrimidine

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	6 <sup>#</sup>	22
Leucovorin	400	1	Q2W	6 <sup>#</sup>	
5-FU	400	1	Q2W	6 <sup>#</sup>	
5-FU	1200*	1-2	Q2W	6 <sup>#</sup>	

\* Continuous infusion for 24 hours #3 (pre-op) + 3 (post-op)

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	6 <sup>#</sup>	19
Leucovorin	200	1	Q2W	6 <sup>#</sup>	
5-FU	2600*	1	Q2W	6 <sup>#</sup>	

\* Continuous infusion for 24 hours #3 (pre-op) + 3 (post-op)

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	130	1	Q3W	6 <sup>#</sup>	23
Capecitabine	1000 PO BID	1-14	Q3W	6 <sup>#</sup>	

#3 (pre-op) + 3 (post-op)

**Oxaliplatin + Fluoropyrimidine**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Docetaxel	50	1	Q2W	8 <sup>#</sup>	8
Oxaliplatin	85	1	Q2W	8 <sup>#</sup>	
Leucovorin	200	1	Q2W	8 <sup>#</sup>	
5-FU	2600*	1	Q2W	8 <sup>#</sup>	

\* Continuous infusion for 24 hours <sup>#</sup>4 (pre-op) + 4 (post-op)

**Other Regimens**
**Cisplatin + Fluorouracil**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	100	1	Q4W	6 <sup>#</sup>	9
5-Fu	800*	1-5	Q4W	6 <sup>#</sup>	

\* Continuous infusion for 24 hours <sup>#</sup>2-3 (pre-op) + 3-4 (post-op)

**Preoperative Chemotherapy**

(Only for adenocarcinoma of the thoracic esophagus or EGJ)

**Cisplatin + Fluorouracil**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	80	1	Q3W	2 <sup>#</sup>	10
5-Fu	1000*	1-4	Q3W	2 <sup>#</sup>	

\* Continuous infusion for 24 hours <sup>#</sup>2(pre-op)

## Definitive Chemoradiation (Non-Surgical) Preferred Regimens

### Cisplatin + Fluoropyrimidine

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	75	1	Q4W	2~4 <sup>#</sup>	11
5-FU	1000*	1-4	Q4W	2~4 <sup>#</sup>	

\* Continuous infusion for 24 hours #2 (RT) + 2

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	30	1	QW	5	42
Capecitabine	800 PO BID	1-5	QW	5	

### Oxaliplatin + Fluoropyrimidine

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	6 <sup>#</sup>	5
Leucovorin	400	1	Q2W	6 <sup>#</sup>	
5-FU	400	1	Q2W	6 <sup>#</sup>	
5-FU	800*	1-2	Q2W	6 <sup>#</sup>	

\* Continuous infusion for 24 hours #3 (RT) + 3 (post-RT)



藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1, 15, 29		For 3 doses	43
Capecitabine	625 PO BID	1-5	QW	5	

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1, 15, 29			3
5-FU	180	1-33			

### Paclitaxel + Carboplatin

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	50	1	QW	5	1
Carboplatin	AUC 2	1	QW	5	

### Other Regimens

#### Taxane + Cisplatin

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	60	1, 8, 15, 22		1	12
Cisplatin	75	1		1	

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Docetaxel	60	1, 22		1	13
Cisplatin	80	1, 22		1	

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Docetaxel	30	1	QW	5	14
Cisplatin	30	1	QW	5	

### Irinotecan + Cisplatin

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Irinotecan	65	1, 8, 22, 29		1	6
Cisplatin	30	1, 8, 22, 29		1	

### Paclitaxel + Fluoropyrimidine

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	45-50	1	QW	5	7
5-FU	300*	1-5	QW	5	

\* Continuous infusion for 24 hours

### Postoperative Chemoradiation (Including EGJ)

#### 5-FU (bolus) + Leucovorin

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Leucovorin	20	1-5	Q4W	1 <sup>st</sup> , 3 <sup>rd</sup> , 4 <sup>th</sup> #	15, 45
5-FU	425	1-5	Q4W	1 <sup>st</sup> , 3 <sup>rd</sup> , 4 <sup>th</sup> #	
Leucovorin	20	1-4, 31-33	Q5W	2 <sup>nd</sup> (with RT)	
5-FU	400	1-4, 31-33	Q5W	2 <sup>nd</sup> (with RT)	

#Cycle 1<sup>st</sup> (pre-RT), and cycle 3<sup>rd</sup>, 4<sup>th</sup> (post-RT)

**5-FU + Leucovorin modifications****1 cycle before + 2 cycle after chemoradiation**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Capecitabine	1000 PO BID	1-14	Q4W	1 + 2	46

**2 cycle before + 4 cycle after chemoradiation**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Leucovorin	400	1	Q2W	2 + 4	
5-FU	400	1	Q2W	2 + 4	
5-FU	1200*	1-2	Q2W	2 + 4	

\*Continuous infusion for 24 hours

**With radiation**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
5-FU	200*	1-5 or 1-7	QW	5 with RT	47

\*Continuous infusion for 24 hours

藥品名 *	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Capecitabine	825 PO BID	1-5 or 1-7	QW	5 with RT	48

## Postoperative Chemotherapy

### Preferred Regimens

#### Nivolumab only after preoperative chemoradiation with R0 resection and residual disease

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Nivolumab	240 mg	1	Q2W	8	50
Followed by					
Nivolumab	480 mg	1	Q4W	For max 1 year	

### Other Recommended Regimens

#### Oxaliplatin + Capecitabine

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	130	1	Q3W		16
Capecitabine	1000 PO BID	1-14	Q3W		

#### Oxaliplatin + Fluorouracil

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W		22
Leucovorin	400	1	Q2W		
5-FU	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

\* Continuous infusion for 24 hours #3 (RT) + 3 (post-RT)

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W		19
Leucovorin	200	1	Q2W		
5-FU	2600*	1-2	Q2W		

\* Continuous infusion for 24 hours #3 (RT) + 3 (post-RT)

## Systemic therapy for metastatic or locally advanced cancer (where local therapy is not indicated)

### First-line Therapy

#### Trastuzumab (with chemotherapy)

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Trastuzumab	8 → 6 mg/kg	1	Q3W		17, 53

#### Trastuzumab (with chemotherapy)

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Trastuzumab	6 → 4 mg/kg	1	Q2W		17

## Preferred Regimens

### ± Nivolumab + Oxaliplatin + Capecitabine

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
± Nivolumab	360 mg	1	Q3W		51
Oxaliplatin	130	1	Q3W		23
Capecitabine	1000 PO BID	1-14	Q3W		

For adenocarcinoma (PD-L1 CPS: 1-4, category 2B; ≥ 5, category 1) or Squamous cell carcinoma.

### ± Nivolumab + Oxaliplatin + 5Fu

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
± Nivolumab	240 mg	1	Q2W		51
Oxaliplatin	85	1	Q2W		22
Leucovorin	400	1	Q2W		
5-FU	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

For adenocarcinoma (PD-L1 CPS: 1-4, category 2B; ≥ 5, category 1) or Squamous cell carcinoma.

\* Continuous infusion for 24 hours

**Pembrolizumab + Oxaliplatin + Capecitabine**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Pembrolizumab	200 mg	1	Q3W	Up to 2 years	52
Oxaliplatin	130	1	Q3W	6	
Capecitabine	1000 PO BID	1-14	Q3W	6	

**Pembrolizumab + Oxaliplatin + 5Fu**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Pembrolizumab	200 mg	1	Q2W	Up to 2 years	52
Oxaliplatin	85	1	Q2W	9	
Leucovorin	400	1	Q2W	9	
5-FU	400	1	Q2W	9	
5-FU	1200*	1-2	Q2W	9	

\* Continuous infusion for 24 hours

**Pembrolizumab + Cisplatin + 5Fu**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Pembrolizumab	200 mg	1	Q3W	Up to 2 years	52
Cisplatin	80	1	Q3W	6	
5-Fu	800*	1-5	Q3W	6	

\* Continuous infusion for 24 hours

**Pembrolizumab + Cisplatin + Capecitabine**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Pembrolizumab	200 mg	1	Q3W	Up to 2 years	52
Cisplatin	80	1	Q3W	6	
Capecitabine	1000 PO BID	1-14	Q3W	6	

### Fluoropyrimidine + Cisplatin

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	100	1	Q4W		18
5-FU	1000*	1-4	Q4W		

\* Continuous infusion for 24 hours

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	50	1	Q2W		19, 20
Leucovorin	200	1	Q2W		
5-FU	2000*	1	Q2W		

\* Continuous infusion for 24 hours

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Cisplatin	80	1	Q3W		21
Capecitabine	1000 PO BID	1-4	Q3W		

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Nivolumab	3 mg/kg	1	Q3W	Max 2 years	54
Ipilimumab	1 mg/kg	1	Q6W		



**Fluoropyrimidine + Oxaliplatin**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W		19
Leucovorin	200	1	Q2W		
5-FU	2600*	1	Q2W		

\* Continuous infusion for 24 hours

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q3W		48
Capecitabine	625 PO BID	1-14	Q3W		

**Other Regimens****Paclitaxel + Cisplatin/Carboplatin**

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	200	1	Q3W		24
Cisplatin	75	2	Q3W		

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	90	1	Q2W		25
Cisplatin	50	1	Q2W		

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	200	1	Q3W		26
Carboplatin	AUC 5	1	Q3W		

### Docetaxel + Cisplatin

藥品名*	劑量 mg/m <sup>2</sup> *	給藥日	頻率	週期	參考文獻
Docetaxel	75-85	1	Q3W		27, 28
Cisplatin	70-75	1	Q3W		

### Fluoropyrimidine

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Leucovorin	400	1	Q2W		20
5-FU	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

\* Continuous infusion for 24 hours

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
5-FU	800*	1-5	Q4W		29

\* Continuous infusion for 24 hours

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Capecitabine	1250 PO BID	1-14	Q3W		30

### Taxane

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Docetaxel	100	1	Q3W		31, 32

藥品名 *	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	250	1	Q3W		33

藥品名 *	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Paclitaxel	80	1	QW		34

### Irinotecan + 5-FU (only for adenocarcinoma)

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Irinotecan	180	1	Q2W		35
Leucovorin	400	1	Q2W		
5-FU	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

\* Continuous infusion for 24 hours

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Irinotecan	80	1	QW	6 + 2(off)#	49
Leucovorin	500	1	QW	6 + 2(off)#	
5-FU	2000*	1	QW	6 + 2(off)#	

\* Continuous infusion for 24 hours #6 weeks treatment, followed by 2 weeks off

### ECF

藥品名 *	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Epirubicin	50	1	Q3W		39
Cisplatin	60	1	Q3W		
5-Fu	200*	1-21	Q3W		

\* Continuous infusion for 24 hours

**ECF modifications**

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Epirubicin	50	1	Q3W		40, 41
Oxaliplatin	130	1	Q3W		
5-Fu	200*	1-21	Q3W		

\* Continuous infusion for 24 hours

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Epirubicin	50	1	Q3W		40, 41
Cisplatin	60	1	Q3W		
Capecitabine	625 PO BID	1-21	Q3W		

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Epirubicin	50	1	Q3W		39
Oxaliplatin	130	1	Q3W		
Capecitabine	625 PO BID	1-21	Q3W		

**DCF modifications**

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Docetaxel	40	1	Q2W		36
Leucovorin	400	1	Q2W		
5-FU	400	1	Q2W		
5-FU	1000*	1-2	Q2W		
Cisplatin	40	3	Q2W		

\* Continuous infusion for 24 hours

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Docetaxel	50	1	Q2W		37
Oxaliplatin	85	1	Q2W		
5-FU	1200*	1-2	Q2W		

\* Continuous infusion for 24 hours

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Docetaxel	75	1	Q3W		38
Carboplatin	AUC 6	2	Q3W		
5-FU	1200*	1-2	Q3W		

\* Continuous infusion for 24 hours

\*三院有個別版本

## 參考文獻

1. Van Hagen P, Hulshof MC, van Lanschot, J.J, et al. Preoperative Chemoradiotherapy for Esophageal or Junctional Cancer. *N Engl J Med* 2012;366:2074-2084.
2. Conroy T, Galais MP, Raoul JL, et al. Definitive chemoradiotherapy with FOLFOX versus fluorouracil and cisplatin in patients with oesophageal cancer (PRODIGE5/ ACCORD17): final results of a randomised, phase 2/3 trial. *Lancet Oncol* 2014;15:305-314.
3. Khushalani NI, Leichman CG, Proulx G, et al. Oxaliplatin in combination with protracted-infusion fluorouracil and radiation: report of a clinical trial for patients with esophageal cancer. *J Clin Oncol* 2002;20:2844-2850.
4. Tepper J, Krasna MJ, Niedzwiecki D, et al. Phase III trial of trimodality therapy with cisplatin, fluorouracil, radiotherapy, and surgery compared with surgery alone for esophageal cancer: CALGB 9781. *J Clin Oncol* 2008;26:1086-1092.
5. Bedenne L, Michel P, Bouche O, et al. Chemoradiation followed by surgery compared with chemoradiation alone in squamous cancer of the esophagus: FFCO 9102. *J Clin Oncol* 2007; 25:1160-1168.
6. Sharma R, Yang GY, Nava HR, et al. A single institution experience with neoadjuvant chemoradiation (CRT) with irinotecan (I) and cisplatin (C) in locally advanced esophageal carcinoma (LAEC) [abstract]. *J Clin Oncol* 2009;27 (Suppl 15):Abstract e15619.
7. Ajani JA, Winter K, Okawara GS, et al. Phase II trial of preoperative chemoradiation in patients with localized gastric adenocarcinoma (RTOG 9904): quality of combined modality therapy and pathologic response. *J Clin Oncol* 2006;24:3953-3958.
8. Al-Batran SE, Hofheinz RD, Pauligk C, et al. Histopathological regression after neoadjuvant docetaxel, oxaliplatin, fluorouracil, and leucovorin versus epirubicin, cisplatin, and fluorouracil or capecitabine in patients with resectable gastric or gastro-oesophageal junction adenocarcinoma (FLOT4-AIO): results from the phase 2 part of a multicentre, open-label, randomised phase 2/3 trial. *Lancet Oncol* 2016;17:1697-1708.
9. Ychou M, Boige V, Pignon J-P, et al. Perioperative chemotherapy compared with surgery alone for resectable

- gastroesophageal adenocarcinoma: an FNCLCC and FFCD multicenter phase III trial. *J Clin Oncol* 2011;29:1715-1721.
10. Alderson D, Langley RE, Nankivell MG, et al. Neoadjuvant chemotherapy for resectable oesophageal and junctional adenocarcinoma: Results from the UK Medical Research Council randomised OEO5 trial (ISRCTN 01852072) [abstract]. *J Clin Oncol* 2015;33 (15\_suppl):Abstract 4002.
  11. Minsky BD, Pajak TF, Ginsberg RJ, et al. INT 0123 (Radiation Therapy Oncology Group 94-05) phase III trial of combined-modality therapy for esophageal cancer: highdose versus standard-dose radiation therapy. *J Clin Oncol* 2002;20:1167-1174.
  12. Urba SG, Orringer MB, Ianettonni M, et al. Concurrent cisplatin, paclitaxel, and radiotherapy as preoperative treatment for patients with locoregional esophageal carcinoma. *Cancer* 003;98:2177-2183.
  13. Li QQ, Liu MZ, Hu YH, et al. Definitive concomitant chemoradiotherapy with docetaxel and cisplatin in squamous esophageal carcinoma. *Dis Esophagus* 2010;23:253-259.
  14. Day FL, Leong T, Ngan S, et al. Phase I trial of docetaxel, cisplatin and concurrent radical radiotherapy in locally advanced oesophageal cancer. *Br J Cancer* 2011;104:265-271.
  15. Smalley SR, Benedetti JK, Haller DG, et al. Updated analysis of SWOG-directed intergroup study 0116: a phase III trial of adjuvant radiochemotherapy versus observation after curative gastric cancer resection. *J Clin Oncol* 2012;30:2327-2333
  16. Noh SH, Park SR, Yang HK, et al. Adjuvant capecitabine plus oxaliplatin for gastric cancer after D2 gastrectomy (CLASSIC): 5-year follow-up of an open-label, randomized phase 3 trial. *Lancet Oncol* 2014; 15:1389-1396.
  17. Bang YJ, Van Cutsem E, Feyereislova A, et al. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet* 2010;376:687-697.
  18. Lorenzen S, Schuster T, Porschen R, et al. Cetuximab plus cisplatin-5-fluorouracil versus cisplatin-5-fluorouracil alone in first-line metastatic squamous cell carcinoma of the esophagus: a randomized phase II study of the Arbeitsgemeinschaft Internistische Onkologie. *Ann Oncol* 2009;20:1667-1673.
  19. Al-Batran S-E, Hartmann JT, Probst S, et al. Phase III trial in metastatic gastroesophageal adenocarcinoma with fluorouracil, leucovorin plus either oxaliplatin or cisplatin: a study of the Arbeitsgemeinschaft Internistische Onkologie. *J*

- Clin Oncol 2008;26:1435-1442.
20. Bouche O, Raoul JL, Bonnetain F, et al. Randomized multicenter phase II trial of a biweekly regimen of fluorouracil and leucovorin (LV5FU2), LV5FU2 plus cisplatin, or LV5FU2 plus irinotecan in patients with previously untreated metastatic gastric cancer: a Federation Francophone de Cancerologie Digestive Group Study--FFCD 9803. *J Clin Oncol* 2004;22:4319-4328.
  21. Kang YK, Kang WK, Shin DB, et al. Capecitabine/cisplatin versus 5-fluorouracil/cisplatin as first-line therapy in patients with advanced gastric cancer: a randomised phase III noninferiority trial. *Ann Oncol* 2009;20:666-673.
  22. Enzinger PC, Burtness B, Hollis D, et al. CALGB 80403/ECOG 1206: A randomized phase II study of three standard chemotherapy regimens (ECF, IC, FOLFOX) plus cetuximab in metastatic esophageal and GE junction cancer [abstract]. *J Clin Oncol* 2010;28 (Suppl 15):Abstract 4006.
  23. Kim GM, Jeung HC, Rha SY, et al. A randomized phase II trial of S-1-oxaliplatin versus capecitabine-oxaliplatin in advanced gastric cancer. *Eur J Cancer* 2012;48:518-526.
  24. Ilson DH, Forastiere A, Arquette M, et al. A phase II trial of paclitaxel and cisplatin in patients with advanced carcinoma of the esophagus. *Cancer J* 2000;6:316-323.
  25. Petrasch S, Welt A, Reinacher A, et al. Chemotherapy with cisplatin and paclitaxel in patients with locally advanced, recurrent or metastatic oesophageal cancer. *Br J Cancer* 1998;78:511-514.
  26. Gadgeel SM, Shields AF, Heilbrun LK, et al. Phase II study of paclitaxel and carboplatin in patients with advanced gastric cancer. *Am J Clin Oncol* 2003;26:37-41.
  27. Ajani JA, Fodor MB, Tjulandin SA, et al. Phase II multi-institutional randomized trial of docetaxel plus cisplatin with or without fluorouracil in patients with untreated, advanced gastric, or gastroesophageal adenocarcinoma. *J Clin Oncol* 2005;23:5660-5667.
  28. Kim JY, Do YR, Park KU, et al. A multi-center phase II study of docetaxel plus cisplatin as first-line therapy in patients with metastatic squamous cell esophageal cancer. *Cancer Chemother Pharmacol* 2010;66:31-36.
  29. Ohtsu A, Shimada Y, Shirao K, et al. Randomized phase III trial of fluorouracil alone versus fluorouracil plus cisplatin



- versus uracil and tegafur plus mitomycin in patients with unresectable, advanced gastric cancer: The Japan Clinical Oncology Group Study (JCOG9205). *J Clin Oncol* 2003;21:54-59.
30. Hong YS, Song SY, Lee SI, et al. A phase II trial of capecitabine in previously untreated patients with advanced and/or metastatic gastric cancer. *Ann Oncol* 2004;15:1344-1347.
  31. Albertsson M, Johansson B, Friesland S, et al. Phase II studies on docetaxel alone every third week, or weekly in combination with gemcitabine in patients with primary locally advanced, metastatic, or recurrent esophageal cancer. *Med Oncol* 2007;24:407-412.
  32. Ford HE, Marshall A, Bridgewater JA, et al. Docetaxel versus active symptom control for refractory oesophagogastric adenocarcinoma (COUGAR-02): an open-label, phase 3 randomised controlled trial. *Lancet Oncol* 2014;15:78-86.
  33. Ajani JA, Ilson DH, Daugherty K, et al. Activity of taxol in patients with squamous cell carcinoma and adenocarcinoma of the esophagus. *J Natl Cancer Inst* 1994;86:1086-1091.
  34. Ilson DH, Wadleigh RG, Leichman LP, Kelsen DP. Paclitaxel given by a weekly 1-h infusion in advanced esophageal cancer. *Ann Oncol* 2007;18:898-902.
  35. Guimbaud R, Louvet C, Ries P, et al. Prospective, randomized, multicenter, phase III study of fluorouracil, leucovorin, and irinotecan versus epirubicin, cisplatin, and capecitabine in advanced gastric adenocarcinoma: A French Intergroup (Fédération Francophone de Cancérologie Digestive, Fédération Nationale des Centres de Lutte Contre le Cancer, and Groupe Coopérateur Multidisciplinaire en ncologie) Study. *J Clin Oncol* 2014;32:3520-3526.
  36. Shah MA, Janjigian YY, Stoller R, et al. Randomized multicenter phase II study of modified docetaxel, cisplatin, and fluorouracil (DCF) versus DCF plus growth factor support in patients with metastatic gastric adenocarcinoma: a study of the US Gastric Cancer Consortium. *J Clin Oncol* 2015;33:3874-3879.
  37. Shankaran V, Mulcahy MF, Hochster HS, et al. Docetaxel, oxaliplatin, and 5-fluorouracil for the treatment of metastatic or unresectable gastric or gastroesophageal junction (GEJ) adenocarcinomas: Preliminary results of a phase II study. *Gastrointestinal Cancers Symposium 2009:Abstract 47*.
  38. Elkerm YM, Elsaid A, AL-Batran S, Pauligk C. Final results of a phase II trial of docetaxel-carboplatin-FU in locally advanced gastric carcinoma [abstract]. Presented at the Gastrointestinal Cancers Symposium 2008. Abstract 38.
  39. Ross P, Nicolson M, Cunningham D, et al. Prospective randomized trial comparing mitomycin, cisplatin, and protracted

- venous-infusion fluorouracil (PVI 5-FU) with epirubicin, cisplatin, and PVI 5-FU in advanced esophagogastric cancer. *J Clin Oncol* 2002;20:1996-2004.
40. Sumpter K, Harper-Wynne C, Cunningham D, et al. Report of two protocol planned interim analyses in a randomised multicentre phase III study comparing capecitabine with fluorouracil and oxaliplatin with cisplatin in patients with advanced oesophagogastric cancer receiving ECF. *Br J Cancer* 2005;92:1976-1983.
  41. Cunningham D, Starling N, Rao S, et al. Capecitabine and oxaliplatin for advanced esophagogastric cancer. *N Engl J Med* 2008;358:36-46.
  42. Le DT, Durham JN, Smith KN, et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* 2017;357:409-413.
  43. Javle MM, Yang G, Nwogu CE, et al. Capecitabine, oxaliplatin and radiotherapy: a phase IB neoadjuvant study for esophageal cancer with gene expression analysis. *Cancer Invest* 2009;27:193-200.
  44. Lee SS, Kim SB, Park SI, et al. Capecitabine and cisplatin chemotherapy (XP) alone or sequentially combined chemoradiotherapy containing XP regimen in patients with three different settings of stage IV esophageal cancer. *Jpn J Clin Oncol* 2007;37:829-835.
  45. Macdonald JS, Smalley SR, Benedetti J, et al. Chemoradiotherapy after surgery compared with surgery alone for adenocarcinoma of the stomach or gastroesophageal junction. *N Engl J Med* 2001;345:725-730.
  46. Jansen EP, Boot H, Saunders MP, et al. A phase I-II study of postoperative capecitabine-based chemoradiotherapy in gastric cancer. *Int J Radiat Oncol Biol Phys* 2007;69:1424-1428.
  47. Leong T, Joon DL, Willis D, et al. Adjuvant chemoradiation for gastric cancer using epirubicin, cisplatin, and 5-fluorouracil before and after three-dimensional conformal radiotherapy with concurrent infusional 5-fluorouracil: a multicenter study of the trans-tasman radiation oncology group. *Int J Radiat Oncol Biol Phys* 2011;79:690-695.
  48. Lee HS, Choi Y, Hur WJ, et al. Pilot study of postoperative adjuvant chemoradiation for advanced gastric cancer: adjuvant 5-FU/cisplatin and chemoradiation with capecitabine. *World J Gastroenterol* 2006;12:603-607.
  49. Wolff K, Wein A, Reulbach U, et al. Weekly high-dose 5-fluorouracil as a 24-h infusion and sodium folinic acid (AIO

- regimen) plus irinotecan in patients with locally advanced nonresectable and metastatic adenocarcinoma or squamous cell carcinoma of the oesophagus: a phase II trial. *Anticancer Drugs* 2009;20:165-173.
50. Kelly R, Ajani J, Kuzdzal J, et al. Adjuvant nivolumab in resected esophageal or gastroesophageal junction cancer following neoadjuvant chemoradiation therapy: first results of the CheckMate 577 study. [abstract]. Presented at the Oral Presentation presented at the ESMO 2020 Annual Meeting; September 19-21, 2020; Virtual Meeting.
51. Moehler M, Shitara K, Garrido M, et al. Nivolumab plus chemotherapy versus chemotherapy as first-line treatment for advanced gastric cancer/gastroesophageal junction cancer/ esophageal adenocarcinoma: first results of the CheckMate 649 study. [abstract]. Presented at the Oral Presentation presented at the ESMO 2020 Annual Meeting; September 19-21, 2020; Virtual Meeting
52. Kato K., Sun JM., M. S, et al. Pembrolizumab plus chemotherapy versus chemotherapy as first-line therapy in patients with advanced esophageal cancer: the phase 3 KEYNOTE-590 study. [abstract]. Presented at the Presented at ESMO Annual Meeting; September 19-21, 2020; Virtual Meeting.
53. Chung HC, et al First-line pembrolizumab/placebo plus trastuzumab and chemotherapy in HER2-positive advanced gastric cancer: KEYNOTE-811. *Future Oncol* 2021;17:491-501.
54. Doki Y, Ajani JA, Kato K, et al. Nivolumab Combination Therapy in Advanced Esophageal Squamous-Cell Carcinoma. *N Engl J Med.* 2022 Feb 3;386(5):449-462.

## 《食道癌放射治療共識》

### 一、治療範圍

1. 食道腫瘤或腫瘤原發部位
2. 淋巴轉移病灶
3. 高風險淋巴轉移範圍

### 二、治療劑量 / 次數

#### ▲輔助性放射治療 (手術後)

##### 1. 總劑量

- (a) 殘餘腫瘤 / 腫瘤原發部位：50Gy(50~50.4Gy) \*
- (b) 高風險範圍：48Gy(45~50.4Gy)
- (c) 照射次數：27 次 (25~30 次)

#### ▲根治性放射治療 (無手術)

##### 1. 總劑量

- (a) 腫瘤：50Gy(50~50.4Gy) \*
- (b) 高風險範圍：48Gy(45~50.4Gy)
- (c) 照射次數：27 次 (25~30 次)

\* 上段食道癌可考慮增加劑量至 60~66 Gy；中下段食道癌可考慮增加劑量至 60 Gy。

#### ▲前導性放射治療 (手術前)

##### 1. 總劑量

- (a) 腫瘤：48Gy(45~50.4Gy)

(b) 高風險範圍：48Gy(45~50.4Gy)

(c) 照射次數：27 次 (25~30 次)

或是

2. 總劑量

(a) 腫瘤：41.4Gy

(b) 高風險範圍：41.4Gy

(c) 照射次數：23 次

### 三、治療方式：

使用強度調控放射治療技術，包含弧形及螺旋放射規畫，可考慮搭配影像導引治療，治療選擇可使用同步照射高與低危險部位的方式或先給予整個照射部位部份劑量照射後，再針對高危險部位加強劑量。

### 四、參考文獻：

1. International Commission on Radiation Units and Measurements. ICRU Report No 50: Prescribing, Recording and Reporting Photon Beam Therapy. Bethesda, MD: ICRU Publications 1993.
2. Cooper JS, GUO MD, Herskovic A, et al. Chemoradiotherapy of locally advanced esophageal cancer: long-term follow-up of a prospective randomized trial (RTOG 85-01) . Radiation Therapy Oncology Group. JAMA 1999;281:1623-1627.
3. International Commission on Radiation Units and Measurements. ICRU Report No 62: Prescribing, Recording and Reporting Photon Beam Therapy. Bethesda, MD: ICRU Publications 1999.
4. Shi XH, Yao W, Liu T. Late course accelerated fractionation in radiotherapy of esophageal carcinoma. Radiother Oncol 1999; 51: 21-26.
5. Minsky BD, Pajak TF, Ginsberg RJ, et al. INT 0123 (Radiation Therapy Oncology Group 94-05) phase III trial of combined-modality therapy for esophageal cancer: high-dose versus standard-dose radiation therapy. J Clin Oncol 2002; 20: 1167-1174.
6. Willett CG, et al. Principles and Practice of Radiation Oncology. 5th edition: Philadelphia: Lippincott Williams & Wilkins; 2007. pp. 1131-1153.

7. P. van Hagen, M.C.C.M. Hulshof, J.J.B. van Lanschot, et al. Preoperative Chemoradiotherapy for Esophageal or Junctional Cancer. *N Engl J Med* 2012; 366:2074-2084.
8. Shapiro J, van Lanschot JJB, Hulshof MCCM, van Hagen P, et al. Neoadjuvant chemoradiotherapy plus surgery versus surgery alone for oesophageal or junctional cancer (CROSS): long-term results of a randomised controlled trial. *Lancet Oncol.* 2015 Sep;16(9):1090-1098.
9. NCCN clinical practice guidelines in oncology for esophageal and esophagogastric junction cancers. Version 5, 2022.