

# 抗癌藥物治療指引

## 肝癌

討論日期

團隊修訂：2019/12/3

核備日期：2020/3/31

# 《肝癌抗癌藥物治療指引》

## Hepatocellular Carcinoma

### Sorafenib (Child-Pugh Class A [category 1] or B)

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Sorafenib	400mg* PO BID				1, 2

\*起始劑量可因病人狀況調整為 200-400 mg PO BID

### Lenvatinib (Child-Pugh Class A only)

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Lenvatinib	8 or 12mg* PO QD				5

\*8 mg/day for BW <60 kg; 12 mg/day for BW ≥60 kg

### Progression on or after Sorafenib

#### Regorafenib (Child-Pugh Class A only [category 1] )

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Regorafenib	160mg PO QD	1-21	Q4W		3

#### Nivolumab (Child-Pugh Class A or B7 only )

藥品名	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Nivolumab	3 mg/kg	1	Q2W		4

#### Ramucirumab (AFP≥400mg/mL only [category 1])

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Ramucirumab	8 mg/kg	1	Q2W		6

#### Pambrolizumab (Child-Pugh Class A [category 2B])

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Pambrolizumab	200 mg	1	Q3W	up to 35	7

#### Atezolizumab + Bevacizumab

藥品名*	劑量 mg/m <sup>2</sup>	給藥日	頻率	週期	參考文獻
Atezolizumab	1200 mg	1	Q3W		8
Bevacizumab	15 mg/kg	1	Q3W		

\*三院有個別版本

## 參考文獻

1. Llovet J, Ricci S, Mazzaferro V, et al. Sorafenib in advanced hepatocellular carcinoma. *New Engl J Med* 2008;359(4):378-390.
2. Cheng A, Kang Y, Chen Z, et al. Efficacy and safety of sorafenib in patients in the Asia Pacific region with advanced hepatocellular carcinoma: a phase III randomized, double-blind, placebo-controlled trial. *Lancet Oncol* 2009;10:25-34.
3. Bruix J, Qin S, Merle P, et al. Regorafenib for patients with hepatocellular carcinoma who progressed on sorafenib treatment (RESORCE): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2017;389:56-66.
4. El-Khoueiry AB, Sangro B, Yau T, et al. Nivolumab in patients with advanced hepatocellular carcinoma (CheckMate 040): an open-label, non-comparative, phase 1/2 dose escalation and expansion trial. *Lancet* 2017;389:2492-2502.
5. Kudo M, Finn RS, Qin S, et al. Lenvatinib versus sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial. *Lancet* 2018;391:1163-1173.
6. Zhu AX, Kang YK, Yen CJ, et al. REACH-2: A randomized, double-blind, placebo-controlled phase 3 study of ramucirumab versus placebo as second-line treatment in patients with advanced hepatocellular carcinoma (HCC) and elevated baseline alpha-fetoprotein (AFP) following first-line sorafenib. *J Clin Oncol* 2018;36:4003.
7. Zhu AX, Finn RS, Edeline J, et al. Pembrolizumab in patients with advanced hepatocellular carcinoma previously treated with sorafenib (KEYNOTE-224): a non-randomised, open-label phase 2 trial. *Lancet Oncol*. 2018 Jul;19(7):940-952.
8. Cheng AL, Qin S, Ikeda M, Galle P, Ducreux M, Zhu A, Kim TY, Kudo M, Breder V, Merle P, Kaseb A, Li D, Verret W, Xu Z, Hernandez S, Liu J, Huang C, Mulla S, Lim HY, Finn R. IMbrave150: Efficacy and safety results from a ph III study evaluating atezolizumab (atezo) + bevacizumab (bev) vs sorafenib (Sor) as first treatment (tx) for patients (pts) with unresectable hepatocellular carcinoma (HCC). *Annals of Oncology*. 2019; 30: ix186~ix187.