

抗癌藥物治療指引

大腸直腸癌

討論日期

團隊修訂：2019/12/4

核備日期：2020/3/31

《大腸直腸癌抗癌藥物治療指引》

Adjuvant Therapy of Colon Cancer

mFOLFOX6

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	12	1-3
Leucovorin	400	1	Q2W	12	
5-FU	400	1	Q2W	12	
5-FU	1200*	1-2	Q2W	12	

* Continuous infusion for 24 hours

FOLFOX4

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	12	8
Leucovorin	200	1	Q2W	12	
5-FU	400	1	Q2W	12	
5-FU	600*	1-2	Q2W	12	

* Continuous infusion for 24 hours

FOLFOX7

藥品名★	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W		11
Leucovorin	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

* Continuous infusion for 24 hours

Capecitabine

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Capecitabine	1250 PO BID	1-14	Q3W	8	4

CapeOx

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	130	1	Q3W	8	5
Capecitabine	1000 PO BID	1-14	Q3W	8	

5 FU + LV

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Leucovorin	500	1, 8, 15, 22, 29, 36	Q8W	4	6
5-FU	500	1, 8, 15, 22, 29, 36	Q8W	4	

sLV5FU2

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Leucovorin	400	1	Q2W	12	7
5-FU	400	1	Q2W	12	
5-FU	1200*	1-2	Q2W	12	

* Continuous infusion for 24 hours

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
UFUR ± Leucovorin	300-350/day PO 50-150 mg PO QD	1-28 1-28	Q4W Q4W	6 6	9

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
TS-1	40 PO BID	1-28	Q6W	4	10

Neoadjuvant Therapy of Colon Cancer

同 1st line therapy of metastatic colon cancer

*三院有個別版本

參考文獻

- Andre T, Boni C, Mounedji-Boudiaf L, et al. Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. *N Engl J Med* 2004;350:2343-2351.
- Cheeseman SL, Joel SP, Chester JD, et al. A 'modified de Gramont' regimen of fluorouracil, alone and with oxaliplatin, for advanced colorectal cancer. *Br J Cancer* 2002;87:393-399.
- Maindrault-Goebel F, deGramont A, Louvet C, et al. Evaluation of oxaliplatin dose intensity in bimonthly leucovorin and 48- hour 5-fluorouracil continuous infusion regimens (FOLFOX) in pretreated metastatic colorectal cancer. *Annals of Oncology* 2000;11:1477-1483.
- Twelves C, Wong A, Nowacki MP, et al. Capecitabine as adjuvant treatment for stage III colon cancer. *N Engl J Med* 2005;352:2696-2704.
- Schmoll HJ, Cartwright T, Taernero J, et al. Phase III trial of capecitabine plus oxaliplatin as adjuvant therapy for stage III colon cancer: a planned safety analysis in 1,864 patients. *J Clin Oncol* 2007;25:102-109.
- Haller DG, Catalano P, Macdonald JS, Mayer RJ. Phase III study of fluorouracil, leucovorin and levamisole in high risk stage II and III colon cancer: final report of Intergroup 0089. *J Clin Oncol* 2005;23:8671-8678.
- Andre T, Louvet C, Maindrault-Goebel F, et al. CPT-11 (irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer. *Eur J Cancer* 1999;35(9):1343-7.
- deGramont A, Figuer A, Seymour M, et al. Leucovorin and Fluorouracil With or Without Oxaliplatin as First-Line Treatment in Advanced Colorectal Cancer. *J Clin Oncol* 2000;18:2938-2947.

9. Sulkes A, Benner SE, Canetta RM. Uracil-florafur: an oral fluoropyrimidine active in colorectal cancer. *J Clin Oncol.* Oct 1998;16(10):3461-3475.
10. Hamaguchi, Tetsuya et al. Capecitabine versus S-1 as adjuvant chemotherapy for patients with stage III colorectal cancer (JCOG0910): an open-label, non-inferiority, randomised, phase 3, multicentre trial. *The Lancet Gastroenterology & Hepatology*, Volume 3 , Issue 1 , 47 – 56.
11. Tezuka T, Hamada C, Ishida H, et al. Phase II clinical study of modified FOLFOX7 (intermittent oxaliplatin administration) plus bevacizumab in patients with unresectable metastatic colorectal cancer-CRAFT study. *Invest New Drugs.* 2013 Oct;31(5):1321-9.

Adjuvant Therapy of Rectal Cancer

Chemotherapy

mFOLFOX6

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	12	1-3
Leucovorin	400	1	Q2W	12	
5-FU	400	1	Q2W	12	
5-FU	1200*	1-2	Q2W	12	

* Continuous infusion for 24 hours

sLV5FU2

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Leucovorin	400	1	Q2W	12	4, 13
5-FU	400	1	Q2W	12	
5-FU	1200*	1-2	Q2W	12	

* Continuous infusion for 24 hours

Capecitabine

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Capecitabine	1250 PO BID	1-14	Q3W	8	5

CapeOx

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	130	1	Q3W	8	6, 7
Capecitabine	1000 PO BID	1-14	Q3W	8	

5 FU + LV

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Leucovorin	500	1, 8, 15, 22, 29, 36	Q8W	4	8
5-FU	500	1, 8, 15, 22, 29, 36	Q8W	4	

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
UFUR	300-350/day PO	1-28	Q4W	6	14
± Leucovorin	50-150 mg PO QD	1-28	Q4W	6	

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
TS-1	40 PO BID	1-28	Q42D	4	15

Chemotherapy + RT

XRT + continuous infusion 5 FU

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
5-FU	225	1-5 or 1-7	Q4W	During XRT	9

XRT + 5 FU/LV

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
5-FU	400	1-4	Q4W	During week 1, 5 of XRT	10
Leucovorin	20	1-4	Q4W		

XRT + Capecitabine

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Capecitabine	825 PO BID	1-5	QW	5	11, 12

XRT + mFOLFOX6

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1			16
Leucovorin	400	1			
5-FU	400	1			
5-FU	1200*	1-2			

* Continuous infusion for 24 hours

Neoadjuvant Therapy of Rectal Cancer

FOLFOX

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	6	16
Leucovorin	400	1	Q2W	6	
5-FU	400	1	Q2W	6	
5-FU	1200*	1-2	Q2W	6	

* Continuous infusion for 24 hours

CapeOx

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	130	1	Q3W	4	17
Capecitabine	1000 PO BID	1-14	Q3W	4	

參考文獻

- Andre T Boni C, Mounedji-Boudiaf L, et al. Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. N Engl J Med 2004;350:2343-2351.
- Cheeseman SL, Joel SP, Chester JD, et al. A 'modified de Gramont' regimen of fluorouracil, alone and with oxaliplatin, for advanced colorectal cancer. Br J Cancer 2002;87:393-399.
- Maindrault-Goebel F, deGramont A, Louvet C, et al. Evaluation of oxaliplatin dose intensity in bimonthly leucovorin and 48- hour 5-fluorouracil continuous infusion regimens (FOLFOX) in pretreated metastatic colorectal cancer. Annals of Oncology 2000;11:1477-1483.

4. Andre T, Louvet C, Maindrault-Goebel F, et al. CPT-11 (irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer. *Eur J Cancer* 1999;35(9):1343-7.
5. Twelves C, Wong A, Nowacki MP, et al. Capecitabine as adjuvant treatment for stage III colon cancer. *N Engl J Med* 2005;352:2696-2704.
6. Schmoll HJ, Cartwright T, Taernero J, et al. Phase III trial of capecitabine plus oxaliplatin as adjuvant therapy for stage III colon cancer: a planned safety analysis in 1,864 patients. *J Clin Oncol* 2007;25:102-109.
7. Haller DG, Tabernero J, Maroun J, et al. Capecitabine Plus Oxaliplatin Compared With Fluorouracil and Folinic Acid As Adjuvant Therapy for Stage III Colon Cancer. *J Clin Oncol* 2011;29:1465-1471.
8. Petrelli N, Douglass Jr HO, Herrare L, et al. The modulation of fluorouracil with leucovorin in metastatic colorectal carcinoma: 40 a prospective randomized phase III trial. *J Clin Oncol* 1989;7:1419-1426.
9. O'Connell MJ, Martenson JA, Wieand HS, et al. Improving adjuvant therapy for rectal cancer by combining protracted-infusion fluorouracil with radiation therapy after curative surgery. *N Engl J Med* 1994;331:502-507.
10. Tepper JE, O'Connell M, Niedzwiecki D, et al. Adjuvant therapy in rectal cancer: analysis of stage, sex, and local control—final report of Intergroup 0114. *J Clin Oncol* 2002;20:1744-1750.
11. O'Connell MJ, Colangelo LH, Beart RW, et al. Capecitabine and oxaliplatin in the preoperative multimodality treatment of rectal cancer: surgical end points from National Surgical Adjuvant Breast and Bowel Project trial R-04. *J Clin Oncol* 2014;32:1927-1934.
12. Hofheinz R, Wenz FK, Post S, et al. Chemoradiotherapy with capecitabine versus fluorouracil for locally advanced rectal cancer: A randomized, multicenter, non-inferiority, phase 3 trial. *Lancet Oncol* 2012;13:579-588.
13. Shikina A, Shinto E, Hashiguchi Y, et al. Differential clinical benefits of 5-fluorouracil-based adjuvant chemotherapy for patients with stage III colorectal cancer according to CD133 expression status. *Jpn J Clin Oncol*. Jan 2014;44(1):42-48.
14. Sulkes A, Benner SE, Canetta RM. Uracil-*ftorafur*: an oral fluoropyrimidine active in colorectal cancer. *J Clin Oncol*. Oct 1998;16(10):3461-3475.
15. Hamaguchi, Tetsuya et al. Capecitabine versus S-1 as adjuvant chemotherapy for patients with stage III colorectal cancer (JCOG0910): an open-label, non-inferiority, randomised, phase 3, multicentre trial. *The Lancet Gastroenterology & Hepatology*, Volume 3 , Issue 1 , 47 – 56.
16. Koizumi M, Yamada T, Shinji S, et al. Feasibility of Neoadjuvant FOLFOX Therapy Without Radiotherapy for Baseline Resectable Rectal Cancer. *In Vivo*. 2018; 32(4): 937–943.
17. Hata T, Takahashi H, Sakai D, et al. Neoadjuvant CapeOx therapy followed by sphincter-preserving surgery for lower rectal cancer. *Surg Today* (2017) 47: 1372.

Chemotherapy for Advanced or Metastatic Colon and Rectal Cancer

First-line therapy

mFOLFOX6 or mFOLFOX7

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	8-12	1, 2, 3, 22
Leucovorin	400	1	Q2W	8-12	
5-FU (optional)	400	1	Q2W	8-12	
5-FU	1200*	1-2	Q2W	8-12	

* Continuous infusion for 24 hours

FOLFOX + Bevacizumab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Bevacizumab	5 mg/kg	1	Q2W		4
Oxaliplatin	85	1	Q2W		
Leucovorin	400	1	Q2W		
5-FU (optional)	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

* Continuous infusion for 24 hours

FOLFOX + Panitumumab (KRAS/NRAS WT gene only)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Panitumumab	6 mg/kg	1	Q2W		5
Oxaliplatin	85	1	Q2W		
Leucovorin	400	1	Q2W		
5-FU (optional)	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

* Continuous infusion for 24 hours

FOLFOX + Cetuximab (KRAS/NRAS WT gene only)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Cetuximab	500 (400→250)	1	Q2W (QW)		6
Oxaliplatin	85	1	Q2W		
Leucovorin	400	1	Q2W		
5-FU (optional)	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

* Continuous infusion for 24 hours

Xelox (Q2W)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	85	1	Q2W	Max 12	26
Capecitabine	1000 PO BID	1-7	Q2W	Max 12	

CapeOx

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	130	1	Q3W	Max 16	7
Capecitabine	1000 PO BID	1-14	Q3W	Max 16	

CapeOx + Bevacizumab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Bevacizumab	7.5 mg/kg	1	Q3W	Max 16	7
Oxaliplatin	130	1	Q3W	Max 16	
Capecitabine	1000 PO BID	1-14	Q3W	Max 16	

FOLFIRI

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Irinotecan	180	1	Q2W		8, 9
Leucovorin	400	1	Q2W		
5-FU (optional)	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

* Continuous infusion for 24 hours

FOLFIRI + Bevacizumab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Bevacizumab	5 mg/kg	1	Q2W		10
Irinotecan	180	1	Q2W		
Leucovorin	400	1	Q2W		
5-FU (optional)	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

* Continuous infusion for 24 hours

FOLFIRI + Cetuximab (KRAS/NRAS WT gene only)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Cetuximab	500 (400→250)	1	Q2W (QW)		11, 12
Irinotecan	180	1	Q2W		
Leucovorin	400	1	Q2W		
5-FU (optional)	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

* Continuous infusion for 24 hours

FOLFIRI + Panitumumab (KRAS/NRAS WT gene only)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Panitumumab	6 mg/kg	1	Q2W		13
Irinotecan	180	1	Q2W		
Leucovorin	400	1	Q2W		
5-FU (optional)	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

* Continuous infusion for 24 hours

Capecitabine

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Capecitabine	1000 (825-1250) PO BID	1-14	Q3W		14

Capecitabine + Bevacizumab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Bevacizumab	7.5 mg/kg	1	Q3W		14
Capecitabine	1000 (825-1250) PO BID	1-14	Q3W		

FOLFOXIRI ± Bevacizumab

藥品名	劑量★ mg/m ²	給藥日	頻率	週期	參考文獻
± Bevacizumab	5 mg/kg	1	Q2W		18, 19
Oxaliplatin	85	1	Q2W		
Irinotecan	180	1	Q2W		
Leucovorin	400	1	Q2W		
5-FU	1600*	1-2	Q2W		

* Continuous infusion for 24 hours

Cetuximab (KRAS/NRAS WT gene only)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Cetuximab	500 (400→250)	1	Q2W (QW)		12, 20

Panitumumab (KRAS/NRAS WT gene only)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Panitumumab	6 mg/kg	1	Q2W		21

Pembrolizumab (MSI-H)

藥品名★	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Pembrolizumab	2 mg/kg	1	Q3W		29

Pembrolizumab (MSI-H)

藥品名★	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Pembrolizumab	200 mg	1	Q3W		23

Nivolumab (MSI-H)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Nivolumab	3 mg/kg	1	Q2W		24

Nivolumab (MSI-H)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Nivolumab	240 mg	1	Q2W		24

Nivolumab + Ipilimumab (MSI-H)

藥品名★	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Nivolumab	3 mg/kg	1	Q3W	4	25
Ipilimumab	1 mg/kg	1	Q3W		
Followed by Nivolumab	3 mg/kg or 240 mg	1	Q2W		

Bolus or Infusional 5 FU/Leucovorin**Roswell Park**

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Leucovorin	500	1, 8, 15, 22, 29, 36	Q8W		15
5-FU	500	1, 8, 15, 22, 29, 36	Q8W		

Simplified Biweekly Infusional 5 FU/LV (sLV5FU2)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Leucovorin	400	1	Q2W		8
5-FU	400	1	Q2W		
5-FU	1200*	1-2	Q2W		

* Continuous infusion for 24 hours

Weekly

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Leucovorin	20	1	QW		16
5-FU	500	1	QW		

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Leucovorin	500	1	QW		17
5-FU	2600*	1	QW		

* Continuous infusion for 24 hours

*三院有個別版本

參考文獻

- deGramont A, Figer A, Seymour M, et al. Leucovorin and Fluorouracil With or Without Oxaliplatin as First-Line Treatment in Advanced Colorectal Cancer. *J Clin Oncol* 2000;18:2938-2947.
- Cheeseman SL, Joel SP, Chester JD, et al. A 'modified de Gramont' regimen of fluorouracil, alone and with oxaliplatin, for advanced colorectal cancer. *Br J Cancer* 2002;87:393-399.
- Maindrault-Goebel F, deGramont A, Louvet C, et al. Evaluation of oxaliplatin dose intensity in bimonthly leucovorin and 48-hour 5-fluorouracil continuous infusion regimens (FOLFOX) in pretreated metastatic colorectal cancer. *Ann Oncol* 2000;11:1477-1483.
- Emmanouilidou C, Sfakiotaki G, Androulakis N, et al. Front-line Bevacizumab in combination with Oxaliplatin, Leucovorin and 5-Fluorouracil (FOLFOX) in patients with metastatic colorectal cancer: a multicenter phase II study. *BMC Cancer* 2007;7:91.
- Douillard JY, Siena S, Cassidy J, et al. Randomized, Phase III Trial of Panitumumab With Infusional Fluorouracil, Leucovorin, and Oxaliplatin (FOLFOX4) Versus FOLFOX4 Alone As First-Line Treatment in Patients With Previously Untreated Metastatic Colorectal Cancer: The PRIME Study. *J Clin Oncol* 2010;28:4697-4705.
- Venook AP, Niedzwiecki D, Lenz H-J, et al. CALGB/SWOG 80405: Phase III trial of irinotecan/5-FU/leucovorin (FOLFIRI) or oxaliplatin/5-FU/leucovorin (mFOLFOX6) with bevacizumab (BV) or cetuximab (CET) for patients (pts) with KRAS wild-type (wt) untreated metastatic adenocarcinoma of the colon or rectum [abstract]. ASCO Meeting Abstracts 2014;32:LBA3.

7. Saltz LB, Clarke S, Diaz-Rubio E, et al. Bevacizumab in Combination With Oxaliplatin-Based Chemotherapy As First-Line Therapy in Metastatic Colorectal Cancer: A Randomized Phase III Study. *J Clin Oncol* 2008;26:2013-2019.
8. Andre T, Louvet C, Maindrault-Goebel F, et al. CPT-11 (Irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer. *Eur J Cancer* 1999;35(9):1343-7.
9. Fuchs CS, Marshall J, Mitchell E, et al. Randomized, Controlled Trial of Irinotecan Plus Infusional, Bolus, or Oral Fluoropyrimidines in First-Line Treatment of Metastatic Colorectal Cancer: Results From the BICC-C Study. *J Clin Oncol* 2007;25:4779-4786.
10. Heinemann V, von Weikersthal LF, Decker T, et al. FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab as first-line treatment for patients with metastatic colorectal cancer (FIRE-3): a randomised, open-label, phase 3 trial. *Lancet Oncol* 2014.
11. Cunningham D, Humblet Y, Siena S, et al. Cetuximab Monotherapy and Cetuximab plus Irinotecan in Irinotecan-Refractory Metastatic Colorectal Cancer. *N Engl J Med* 2004;351:337-345.
12. Martin-Martorell P, Rosello' S, Rodriguez-Braun E, et al. Biweekly cetuximab and irinotecan in advanced colorectal cancer 50 patients progressing after at least one previous line of chemotherapy: results of a phase II single institution trial. *Br J Cancer* 2008;99:455-458.
13. Peeters M, Prince TJ, Cervantes A, et al. Randomized Phase III Study of Panitumumab With Fluorouracil, Leucovorin, and Irinotecan (FOLFIRI) Compared With FOLFIRI Alone As Second-Line Treatment in Patients With Metastatic Colorectal Cancer. *J Clin Oncol* 2010;28:4706-4713.
14. Cunningham D, Lang I, Marcuello E, et al. Bevacizumab plus capecitabine versus capecitabine alone in elderly patients with previously untreated metastatic colorectal cancer (AVEX): an open-label, randomised phase 3 trial. *Lancet Oncol* 2013;14:1077-1085.
15. Wolmark N, Rockette H, Fisher B, et al. The benefit of leucovorin-modulated fluorouracil as postoperative adjuvant therapy for primary colon cancer: results from National Surgical Adjuvant Breast and Bowel Project protocol C-03. *J Clin Oncol* 1993;11:1879-1887.
16. Jäger E, Heike M, Bemhard H, et al. Weekly high-dose leucovorin versus low-dose leucovorin combined with fluorouracil in advanced colorectal cancer: results of a randomized multicenter trial. *J Clin Oncol* 1996;14:2274-2279.
17. Douillard JY, Cunningham D, Roth AD, et al. Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: a multicentre randomised trial. *The Lancet* 2000;355:1041-47.
18. Falcone A, Ricci S, Brunetti I, et al. Phase III Trial of Infusional Fluorouracil, Leucovorin, Oxaliplatin, and Irinotecan (FOLFOXIRI) Compared With Infusional Fluorouracil, Leucovorin, and Irinotecan (FOLFIRI) As First-Line Treatment for Metastatic Colorectal Cancer: The Gruppo Oncologico Nord Ovest. *J Clin Oncol* 2007;25(13):1670-1676.

19. Cremolini C, Loupakis F, Antoniotti C, et al. FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as first-line treatment of patients with metastatic colorectal cancer: updated overall survival and molecular subgroup analyses of the open-label, phase 3 TRIBE study. *Lancet Oncol* 2015;16:1306-1315.
20. Van Cutsem E, Tejpar S, Vanbekevoort D, et al. Intrapatient Cetuximab Dose Escalation in Metastatic Colorectal Cancer According to the Grade of Early Skin Reactions: The Randomized EVEREST Study. *J Clin Oncol* 2012;30:2861-2868.
21. Van Cutsem E, Peeters M, Siena S, et al. Open-Label Phase III Trial of Panitumumab Plus Best Supportive Care Compared With Best Supportive Care Alone in Patients With Chemotherapy-Refractory Metastatic Colorectal Cancer. *J Clin Oncol* 2007;25:1658-1664.
22. Hochster HS, Grothey A, Hart L, et al. Improved time to treatment failure with an intermittent oxaliplatin strategy: results of CONCePT. *Ann Oncol* 2014;25:1172-1178.
23. Le DT, Uram JN, Wang H, et al. PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med*. Jun 25 2015;372(26):2509-2520.
24. Overman MJ KS, McDermott RS, et al. Nivolumab {+/-} ipilimumab in treatment of patients with metastatic colorectal cancer (mCRC) with and without high microsatellite instability (MSI-H): CheckMate-142 interim results [abstract]. *ASCO Meeting Abstracts* 2016;34:3501. 2016.
25. Overman MJ, Lonardi S, Wong K, et al. Durable Clinical Benefit With Nivolumab Plus Ipilimumab in DNA Mismatch Repair–Deficient/Microsatellite Instability–High Metastatic Colorectal Cancer. *J Clin Oncol* 2018;36:773-779.
26. Grande C, Quintero G, Candamio S, et al. Biweekly XELOX (capecitabine and oxaliplatin) as first-line treatment in elderly patients with metastatic colorectal cancer. *Journal of Geriatric Oncology* 2013;4: 114-121.