

《 肝癌放射治療共識 》

一、治療範圍

1. 肝腫瘤
2. 栓塞肝門靜脈

二、治療劑量 / 次數 (依正常肝臟劑量調整)

▲ 常規治療：

30-66Gy

次數 10 次 -33 次

▲ 立體定位高劑量分次治療

1. 分次劑量：6-20Gy
2. 治療次數：3-6 次

三、治療方式：

使用立體順型與強度調控放射治療技術，包含弧形及螺旋放射規畫，可考慮搭配影像導引治療，治療選擇可使用同步照射高與低危險部位的方式或先給予整個照射部位部份劑量照射後，再針對高危險部位加強劑量。

四、參考文獻：

1. NCCN clinical practice guidelines in oncology-Hepatobiliary cancers. Version 3.2019.
2. Yoon HI, et al. Improved oncologic outcomes with image-guided intensity-modulated radiation therapy using helical tomotherapy in locally advanced hepatocellular carcinoma. J Cancer Res Clin Oncol. 2014 Sep;140(9):1595-605.
3. Cardenes HR, et al. Phase I feasibility trial of stereotactic body radiation therapy for primary hepatocellular carcinoma. Clin Transl Oncol 2010; 12: 218-225.
4. Tse RV, et al. Phase I study of individualized stereotactic body radiotherapy for hepatocellular carcinoma and intrahepatic cholangiocarcinoma. J Clin Oncol 2008; 26: 657-664.

5. Bentzen SM, Constine LS, Deasy JO, et al. Quantitative Analyses of Normal Tissue Effects in the Clinic (QUANTEC): an introduction to the scientific issues. *International journal of radiation oncology, biology, physics* 2010; 76(3 Suppl): S3-9
6. Nitin Ohri, Laura A. Dawson, Sunil Krishnan, et al. Radiotherapy for Hepatocellular Carcinoma: New Indications and Directions for Future Study. *J Natl Cancer Inst.* 2016 Jul 4;108(9).
7. Chen SW, Lin LC, Kuo YC, et al. Phase 2 study of combined sorafenib and radiation therapy in patients with advanced hepatocellular carcinoma. *Int J Radiat Oncol Biol Phys.* 2014 Apr 1;88(5):1041-7.
8. Lee HL, Tsai JT, Chen CY, et al. Effectiveness of stereotactic ablative radiotherapy in patients with advanced hepatocellular carcinoma unsuitable for transarterial chemoembolization. *Ther Adv Med Oncol.* 2019 Dec 4;11:1758835919889002.