

《 淋巴瘤抗癌藥物治療指引 》

Hodgkin Lymphoma (Age ≥ 18 years)

Classical Hodgkin Lymphoma

ABVD (Doxorubicin, Bleomycin, Vinblastine, Dacarbazine) ± ISRT

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Doxorubicin	25	1, 15	Q4W	4	1-4
Bleomycin	10 unit/m ²	1, 15	Q4W	4	
Vinblastine	6	1, 15	Q4W	4	
Dacarbazine	375	1, 15	Q4W	4	

Escalated BEACOPP (Bleomycin, Etoposide, Doxorubicin, Cyclophosphamide, Vincristine, Procarbazine, Prednisone) followed by

ABVD with ISRT10

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Bleomycin	10 unit/m ²	8	Q2W	2 + 2	5-6
Etoposide	200	1-3	Q2W	2 + 2	
Doxorubicin	35	1	Q2W	2 + 2	
Cyclophosphamide	1200	1	Q2W	2 + 2	
Vincristine	1.4	8	Q2W	2 + 2	
Procarbazine	100 QHS	1-7	Q2W	2 + 2	
Prednisone	40 PO QD	1-14	Q2W	2 + 2	

Brentuximab vedotin+AVD (doxorubicin+ vinblastine+ dacarbazine)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Brentuximab vedotin	1.2 mg/kg	1, 15	Q4W	6	7
Doxorubicin	25	1, 15	Q4W	6	
Vinblastine	6	1, 15	Q4W	6	
Dacarbazine	375	1, 15	Q4W	6	

*三院有個別版本

參考文獻

1. Eich HT, Diehl V, Gorgen H, et al. Intensified chemotherapy and dose-reduced involved-field radiotherapy in patients with early unfavorable Hodgkin's lymphoma: final analysis of the German Hodgkin Study Group HD 11 trial. *J Clin Oncol* 2010;28:4199-4206.
2. Engert A, Plutschow A, Eich HT, et al. Reduced treatment intensity in patients with early-stage Hodgkin's lymphoma. *N Engl J Med* 2010;363:640-652.
3. Raemaekers JM, André MP, Federico M, et al. Omitting radiotherapy in early positron emission tomography-negative stage I/II Hodgkin lymphoma is associated with an increased risk of early relapse: clinical results of the preplanned interim analysis of the randomized EORTC/LYSA/FIL H10 trial. *J Clin Oncol* 2014;32:1188-1194.
4. Radford J, Illidge T, Counsell N, et al. Results of a trial of PET-directed therapy for early-stage Hodgkin's lymphoma. *N Engl J Med* 2015;372:1598-1607.
5. Twelves C, Wong A, Nowacki MP, et al. Capecitabine as adjuvant treatment for stage III colon cancer. *N Engl J Med* 2005;352:2696-2704.
6. Engert A, Haverkamp H, Cobe C, et al. Reduced-intensity chemotherapy and PET-guided radiotherapy in patients with advanced stage Hodgkin's lymphoma (HD 15 trial): a randomized, open-label, phase 3 non-inferiority trial. *Lancet* 2012; 379(9828): 1791-1799

6. von Tresckow B, Plutschow A, Fuches M, et al. Dose-intensification in early unfavorable Hodgkin's lymphoma: final analysis of the German Hodgkin Study Group HD14 trial. J Clin Oncol 2012; 30: 907-913.
7. Connors JM, Jurczak W, Straus DJ, et al. Brentuximab Vedotin with Chemotherapy for Stage III or IV Hodgkin's Lymphoma. New England Journal of Medicine. 2018;378(4):331-344.

Nodular Lymphocyte-Predominant Hodgkin Lymphoma

ABVD (Doxorubicin, Bleomycin, Vinblastine, Dacarbazine) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Doxorubicin	25	1, 15	Q4W		1, 2
Bleomycin	10 unit/m ²	1, 15	Q4W		
Vinblastine	6	1, 15	Q4W		
Dacarbazine	375	1, 15	Q4W		
± Rituximab	375	1	Q4W		

CHOP (Cyclophosphamide, Doxorubicin, Vincristine, Prednisone) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Cyclophosphamide	750	1	Q3W		3
Doxorubicin	50	1	Q3W		
Vincristine	1.4	1	Q3W		
Prednisone	40	1-5	Q3W		
± Rituximab	375	1	Q3W		

CVP (Cyclophosphamide, Vinblastine, Prednisone) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Cyclophosphamide	500	1	Q2-3W		4
Vinblastine	6	1, 8	Q2-3W		
Prednisone	40	1-7	Q2-3W		
± Rituximab	375	1	Q2-3W		

Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	QW		5-9

參考文獻

1. Savage KJ, Skinnider B, Al-Mansour M, et al. Treating limited stage nodular lymphocyte predominant Hodgkin lymphoma similarly to classical Hodgkin lymphoma with ABVD may improve outcome. *Blood* 2011;118:4585-4590.
2. Canellos GP, Mauch P. What is the appropriate systemic chemotherapy for lymphocyte-predominant Hodgkin's Lymphoma? *J Clin Oncol* 2010;28:e8.
3. Fanale MA, Lai C-M, McLaughlin P, et al. Outcomes of Nodular Lymphocyte Predominant Hodgkin's Lymphoma (NLPHL) Patients Treated with R-CHOP. *ASH Annual Meeting Abstracts* 2010;116:2812.
4. Shankar A, Hall GW, Gorde-Grosjean S, et al. Treatment outcome after low intensity chemotherapy [CVP] in children and adolescents with early stage nodular lymphocyte predominant Hodgkin's lymphoma - an Anglo-French collaborative report. *Eur J Cancer* 2012;48:1700-1706.
5. Advani RH, Hoppe RT. How I treat nodular lymphocyte predominant Hodgkin lymphoma. *Blood* 2013;122:4182-4188.
6. Advani RH, Horning SJ, Hoppe RT, et al. Mature results of a phase II study of rituximab therapy for nodular lymphocyte-predominant Hodgkin lymphoma. *J Clin Oncol* 2014;32:912-918.

7. Schulz H, Rehwald U, Morschhauser F, et al. Rituximab in relapsed lymphocyte-predominant Hodgkin lymphoma: long-term results of a phase 2 trial by the German Hodgkin Lymphoma Study Group (GHSg). *Blood* 2008;111(1):109-111.
8. Eichenauer DA, Fuchs M, Plutschow A, et al. Phase 2 study of rituximab in newly diagnosed stage IA nodular lymphocyte-predominant Hodgkin lymphoma: a report from the German Hodgkin Study Group. *Blood* 2011;118:4363-4365.
9. Eichenauer DA, Plutschow A, Fuchs M, et al. Long-Term Course of Patients With Stage IA Nodular Lymphocyte-Predominant Hodgkin Lymphoma: A Report From the German Hodgkin Study Group. *J Clin Oncol* 2015;33:2857-2862.

Systemic therapy for relapsed or refractory disease

Second-Line or Subsequent Therapy Options

CHL

DHAP (Dexamethasone, Cisplatin, high-dose Cytarabine)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Dexamethasone	40 mg QD	1-4	Q3-4W		1, 2
Cisplatin	100	1	Q3-4W		
Cytarabine	2000 Q12H	2	Q3-4W		

ESHAP (Etoposide, Methylprednisolone, Cisplatin, high-dose Cytarabine)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Etoposide	40	1-4	Q3-4W		3, 4, 5
Methylprednisolone	500	1-4	Q3-4W		
Cisplatin	25	1-4	Q3-4W		
Cytarabine	2000	5	Q3-4W		

Gemcitabine/Bendamustine/Vinorelbine

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Gemcitabine	800	1, 4	Q3W	4	22
Bendamustine	100	2, 3	Q3W	4	
Vinorelbine	20	1	Q3W	4	
Prednisolone	100 mg PO	1-4	Q3W	4	

GCD (Gemcitabine, Carboplatin, Dexamethasone)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Gemcitabine	1000	1, 8	Q3W		6, 7
Carboplatin	AUC 5	1	Q3W		
Dexamethasone	40 mg	1-4	Q3W		

GVD (Gemcitabine, Vinorelbine, Lipo-Doxorubicin)

1. For transplant- naïve patients

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Gemcitabine	1000	1, 8	Q3W		8
Vinorelbine	20	1, 8	Q3W		
Lipo-Doxorubicin	15	1, 8	Q3W		

2. For post-transplant patients

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Gemcitabine	800	1, 8	Q3W		8
Vinorelbine	15	1, 8	Q3W		
Lipo-Doxorubicin	10	1, 8	Q3W		

ICE (Ifosfamide, Carboplatin, Etoposide)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Etoposide	100	1-3	Q3W		9, 10
Carboplatin	AUC 5	2	Q3W		
Ifosfamide	5000	2	Q3W		

IGEV (Ifosfamide, Gemcitabine, Vinorelbine, Prednisolone)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Ifosfamide	2000	1-4	Q3W		11
Gemcitabine	800	1, 4	Q3W		
Vinorelbine	20	1	Q3W		
Prednisolone	100 mg	1-4	Q3W		

Mini-BEAN (Carmustine, Cytarabine, Etoposide, Mephalan)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Carmustine	60	1	Q4-6W		12, 13
Cytarabine	100	2-5	Q4-6W		
Etoposide	75	2-5	Q4-6W		
Mephalan	30	6	Q4-6W		

MINE (Etoposide, Ifosfamide, Mesna, Mitoxantrone)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Mesna	1300	1-3	Q3-4W		14
Ifosfamide	1300	1-3	Q3-4W		
Mitoxantrone	8	1	Q3-4W		
Etoposide	65	1-3	Q3-4W		

Brentuximab vedotin (only for CHL)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Brentuximab vedotin*	1.8 mg/kg	1	Q3W		15

*alone or in combination with the second-line regimens below

Brentuximab vedotin + Bendamustine

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Brentuximab vedotin*	1.8 mg/kg	1	Q3W		23
Bendamustine	90 (70-90)	1, 2	Q3W		

Brentuximab vedotin + Nivolumab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Brentuximab vedotin*	1.8 mg/kg	1	Q3W		24
Nivolumab	3 mg/kg	8 → 1	Q3W	1 st → 2 nd -4 th	

Additional Therapy Options: (only for CHL)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Bendamustine	120	1, 2	Q4W		16

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Everolimus	10 mg PO QD				17

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Lenalidomide	25 mg PO QD	1-21	Q4W		18

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Nivolumab	3 mg/kg	1	Q2W		19, 20

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Pembrolizumab	10 mg/kg	1	Q2W		21

NLPHL**DHAP (Dexamethasone, Cisplatin, high-dose Cytarabine) ± Rituximab**

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1			1, 2
Dexamethasone	40 mg QD	1-4	Q3-4W		
Cisplatin	100	1	Q3-4W		
Cytarabine	2000 Q12H	2	Q3-4W		

ESHAP (Etoposide, Methylprednisolone, Cisplatin, high-dose Cytarabine) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1			3, 4, 5
Etoposide	40	1-4	Q3-4W		
Methylprednisolone	500	1-4	Q3-4W		
Cisplatin	25	1-4	Q3-4W		
Cytarabine	2000	5	Q3-4W		

ICE (Ifosfamide, Carboplatin, Etoposide) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1			9, 10
Etoposide	100	1-3	Q3W		
Carboplatin	AUC 5	2	Q3W		
Ifosfamide	5000	2	Q3W		

IGEV (Ifosfamide, Gemcitabine, Vinorelbine, Prednisolone) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1			11
Ifosfamide	2000	1-4	Q3W		
Gemcitabine	800	1, 4	Q3W		
Vinorelbine	20	1	Q3W		
Prednisolone	100 mg	1-4	Q3W		

參考文獻

1. Josting A, Rudolph C, Reiser M, et al. Time-intensified dexamethasone/cisplatin/cytarabine: an effective salvage therapy with low toxicity in patients with relapsed and refractory Hodgkin's disease. *Ann Oncol* 2002;13(10):1628-1635.
2. Abali H, Urün Y, Oksüzoğlu B, Budakoğlu B, et al. Comparison of ICE (ifosfamide-carboplatinetoposide) versus DHAP (cytosine arabinoside-cisplatin-dexamethasone) as salvage chemotherapy in patients with relapsed or refractory lymphoma. *Cancer Invest* 2008;26(4):401-406.
3. Aparicio J, Segura A, Garcera S, et al. ESHAP is an active regimen for relapsing Hodgkin's disease. *Ann Oncol* 1999;10(5):593-595.
4. Fernández de Larrea C, Martínez C, et al. Salvage chemotherapy with alternating MINEESHAP regimen in relapsed or refractory Hodgkin's lymphoma followed by autologous stem cell transplantation. *Ann Oncol* 2010;21(6):1211-1216.
5. Labrador J, Cabrero-Calvo M, Perez-Lopez E, et al. ESHAP as salvage therapy for relapsed or refractory Hodgkin's lymphoma. *Ann Hematol* 2014;93:1745-1753.
6. Crump M, Kuruvilla J, Couban S, et al. Randomized comparison of gemcitabine, dexamethasone, and cisplatin versus dexamethasone, cytarabine, and cisplatin chemotherapy before autologous stem-cell transplantation for relapsed and refractory aggressive lymphomas: NCIC-CTG LY.12. *J Clin Oncol* 2014;32:3490-3496.
7. Gopal AK, Press OW, Shustov AR, et al. Efficacy and safety of gemcitabine, carboplatin, dexamethasone, and rituximab in patients with relapsed/refractory lymphoma: a prospective multicenter phase II study by Puget Sound Oncology Consortium. *Leuk Lymphoma* 2010;51:1523-1529.
8. Bartlett N, Niedzwiecki D, Johnson J, et al. Gemcitabine, vinorelbine, and pegylated liposomal doxorubicin (GVD), a salvage regimen in relapsed Hodgkin's lymphoma: CALGB 59804. *Ann Oncol* 2007;18(6):1071-1079.
9. Moskowitz CH, Nimer SD, Zelenetz AD, et al. A 2-step comprehensive high-dose chemoradiotherapy second-line program for relapsed and refractory Hodgkin disease: analysis by intent to treat and development of a prognostic model. *Blood* 2001;97(3):616-623.
10. Abali H, Urün Y, Oksüzoğlu B, Budakoğlu B, et al. Comparison of ICE (ifosfamide-carboplatinetoposide) versus DHAP (cytosine arabinoside-cisplatin-dexamethasone) as salvage chemotherapy in patients with relapsed or refractory lymphoma.

- Cancer Invest 2008;26(4):401-406.
11. Santoro A, Magagnoli M, Spina M, et al. Ifosfamide, gemcitabine, and vinorelbine: a new induction regimen for refractory and relapsed Hodgkin's lymphoma. *Haematologica* 2007;92(1):35-41.
 12. Colwill R, Crump M, Couture F, et al. Mini-BEAM as salvage therapy for relapsed or refractory Hodgkin's disease before intensive therapy and autologous bone marrow transplantation. *J Clin Oncol* 1995;13:396-402.
 13. Martín A, Fernández-Jiménez MC, Caballero MD, et al. Long-term follow-up in patients treated with Mini-BEAM as salvage therapy for relapsed or refractory Hodgkin's disease. *Br J Haematol* 2001;113(1):161-171.
 14. Rodriguez MA, Cabanillas FC, Hagemeister FB, et al. A phase II trial of mesna/ifosfamide, mitoxantrone and etoposide for refractory lymphomas. *Ann Oncol* 1995;6(6):609-611.
 15. Younes A, Gopal AK, Smith SE, et al. Results of a pivotal phase II study of brentuximab vedotin for patients with relapsed or refractory Hodgkin's lymphoma. *J Clin Oncol* 2012;30:2183-2189.
 16. Moskowitz AJ, Hamlin PA, Perales M-A, et al. Phase II study of bendamustine in relapsed and refractory Hodgkin lymphoma. *J Clin Oncol* 2013;31:456-460.
 17. Johnston PB, Inwards DJ, Colgan JP, et al; A Phase II trial of the oral mTOR inhibitor everolimus in relapsed Hodgkin lymphoma. *Am J Hematol*. 2010;85(5):320-4.
 18. Fehniger TA, Larson S, Trinkaus K, et al; A phase 2 multicenter study of lenalidomide in relapsed or refractory classical Hodgkin lymphoma. *Blood* 2011;118(19):5119-25.
 19. Ansell SM, Lesokhin AM, Borrello I, et al. PD-1 Blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma. *N Engl J Med* 2015;372:311-9.
 20. Timmerman J, Armand P, Lesokhin AM, et al. Nivolumab in patients with relapsed or refractory lymphoid malignancies and classical Hodgkin lymphoma: Updated results of a phase 1 study (CA 209-039) [abstract]. *Hematol Oncol* 2015;33:Abstract 010.
 21. Moskowitz CH, Ribrag V, Michot J-M, et al. PD-1 blockade with the monoclonal antibody pembrolizumab (MK-3475) in patients with classical Hodgkin lymphoma after brentuximab vedotin failure: preliminary results from a phase 1b study

- (KEYNOTE-013) [abstract]. Blood 2014;124:Abstract 290.
22. Santoro A, Mazza R, Pulsoni A, et al. Bendamustine in Combination With Gemcitabine and Vinorelbine Is an Effective Regimen As Induction Chemotherapy Before Autologous Stem-Cell Transplantation for Relapsed or Refractory Hodgkin Lymphoma: Final Results of a Multicenter Phase II Study. J Clin Oncol 2016;34:3293-3299.
 23. O'Connor OA, Lue JK, et al. Brentuximab vedotin plus bendamustine in relapsed or refractory Hodgkin's lymphoma: an international, multicentre, single-arm, phase 1-2 trial. Lancet Oncol 2018; 19: 257-66.
 24. Herrera AF, Moskowitz AJ, et al. Interim results of brentuximab vedotin in combination with nivolumab in patients with relapsed or refractory Hodgkin lymphoma. Blood (2018) 131 (11): 1183-1194.

Hodgkin Lymphoma (Age > 60 years)

A(B)VD ± ISRT

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Doxorubicin	25	1, 15	Q4W	#	1-4, 7
Bleomycin*	10 unit/m ²	1, 15	Q4W	#	
Vinblastine	6	1, 15	Q4W	#	
Dacarbazine	375	1, 15	Q4W	#	

* Bleomycin should be used with caution as it may not be tolerated in older adults.

A(B)VD (2 cycles) followed by AVD (4 cycles), if PET scan is negative after 2 cycles of ABVD.

If stage I-II unfavorable, consider a total of 4 cycles

CHOP ± ISRT

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Cyclophosphamide	750	1	Q3W	*	5
Doxorubicin	50	1	Q3W	*	
Vincristine	1.4	1	Q3W	*	
Prednisone	40	1-5	Q3W	*	

* Stage I-II favorable disease: 4; Stage I-II favorable or III-IV: 6

PVAG (Prednisone, Vinblastine, Doxorubicin, Gemcitabine)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Prednisone	40	1-5	Q3W	6	6
Vinblastine	6	1	Q3W	6	
Doxorubicin	50	1	Q3W	6	
Gemcitabine	1000	1	Q3W	6	

參考文獻

1. Engert A, Plutschow A, Eich HT, et al. Reduced treatment intensity in patients with early-stage Hodgkin's lymphoma. *N Engl J Med* 2010;363:640-652.
2. Stamatoullas A, Brice P, Bouabdallah R, et al. Outcome of patients older than 60 years with classical Hodgkin lymphoma treated with front line ABVD chemotherapy: frequent pulmonary events suggest limiting the use of bleomycin in the elderly. *Br J Haematol* 2015;170:179-184.
3. Behringer K, Goergen H, Hitz F, et al. Omission of dacarbazine or bleomycin, or both, from the ABVD regimen in treatment of early-stage favourable Hodgkin's lymphoma (GHSG HD13): an open-label, randomised, non-inferiority trial. *Lancet* 2015;385:1418-1427.
4. Johnson P, Federico M, Fossa A, et al. Response-adapted therapy based on interim FDG-PET scans in advanced Hodgkin lymphoma: first analysis of the safety of de-escalation and efficacy of escalation in the international RATHL study (CRUK/07/033) [abstract]. *Hematol Oncol* 2015;33 (Suppl S1):Abstract 008.
5. Kolstad A, Nome O, Delabie J, et al. Standard CHOP-21 as first line therapy for elderly patients with Hodgkin's lymphoma. *Leuk Lymphoma* 2007;48:570-576.
6. Boll B, Bredenfeld H, Gorgen H, et al. Phase 2 study of PVAG (prednisone, vinblastine, doxorubicin, gemcitabine) in elderly patients with early unfavorable or advanced stage Hodgkin lymphoma. *Blood* 2011;118:6292-6298.
7. Johnson P, Federico M, Fossa A, et al. Response-adapted therapy based on interim FDG-PET scans in advanced Hodgkin lymphoma: first analysis of the safety of de-escalation and efficacy of escalation in the international RATHL study (CRUK/07/033) [abstract]. *Hematol Oncol* 2015;33 (Suppl S1): Abstract 008.

Non-Hodgkin's Lymphoma

Diffuse Large B-Cell Lymphoma

First-line Therapy

RCHOP (Rituximab, Cyclophosphamide, Doxorubicin, Vincristine, Prednisone)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	6	1-3
Cyclophosphamide	750	1	Q3W	6	
Doxorubicin	50	1	Q3W	6	
Vincristine	1.4	1	Q3W	6	
Prednisone	100 mg	1-5	Q3W	6	

Dose-dense RCHOP 14

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q2W	6	4
Cyclophosphamide	750	1	Q2W	6	
Doxorubicin	50	1	Q2W	6	
Vincristine	1.4	1	Q2W	6	
Prednisone	100 mg	1-5	Q2W	6	

Dose-adjusted EPOCH + Rituximab**(Etoposide, Prednisone, Vincristine, Cyclophosphamide, Doxorubicin) + Rituximab**

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	6-8	5, 6
Etoposide	50	1-4	Q3W	6-8	
Doxorubicin	10	1-4	Q3W	6-8	
Vincristine	0.4	1-4	Q3W	6-8	
Cyclophosphamide	750	5	Q3W	6-8	
Prednisone	60	1-5	Q3W	6-8	

First-line Therapy for Patients with Poor Left Ventricular Function**CDOP (Cyclophosphamide, Lipo-Doxorubicin, Vincristine, Prednisone) + Rituximab**

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	6-8	7, 8
Cyclophosphamide	750	1	Q3W	6-8	
Lipo-Doxorubicin	30	1	Q3W	6-8	
Vincristine	1.4	1	Q3W	6-8	
Prednisone	60	1-5	Q3W	6-8	

RGCVP (Rituximab, Gemcitabine, Cyclophosphamide, Vincristine, Prednisone)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	6	10
Cyclophosphamide	750	1	Q3W	6	
Gemcitabine	750-1000	1,8	Q3W	6	
Vincristine	1.4	1	Q3W	6	
Prednisone	100 mg	1-5	Q3W	6	

DA-EPOCH (Etoposide, Prednisone, Vincristine, Cyclophosphamide, Doxorubicin) + Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	6-8	5, 6
Etoposide	50	1-4	Q3W	6-8	
Doxorubicin	10	1-4	Q3W	6-8	
Vincristine	0.4	1-4	Q3W	6-8	
Cyclophosphamide	750	5	Q3W	6-8	
Prednisone	60	1-5	Q3W	6-8	

RCEOP (Rituximab, Cyclophosphamide, Etoposide, Vincristine, Prednisone)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	*	9
Cyclophosphamide	750	1	Q3W	*	
Etoposide	50	1	Q3W	*	
Etoposide	100 PO	2, 3	Q3W	*	
Vincristine	1.4	1	Q3W	*	
Prednisone	100 mg	1-5	Q3W	*	

*limited stage: 3-4, advanced stage: 6

TREC (Rituximab, Bendamustine, Etoposide, Carboplatin)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	2	15
Bendamustine	90-120	1-2	Q3W	2	
Etoposide	100	1-3	Q3W	2	
Carboplatin	AUC 5	1	Q3W	2	

Patients >80years of age with comorbidities**R-mini-CHOP**

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	6	11
Cyclophosphamide	400	1	Q3W	6	
Doxorubicin	25	1	Q3W	6	
Vincristine	1 mg	1	Q3W	6	
Prednisone	40	1-5	Q3W	6	

R-COP

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W		24-25
Cyclophosphamide	750	1	Q3W		
Vincristine	1.4 mg	1	Q3W		
Prednisone	100	1-5	Q3W		

RGCVP (Rituximab, Gemcitabine, Cyclophosphamide, Vincristine, Prednisolone)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	6	10
Cyclophosphamide	750	1	Q3W	6	
Gemcitabine	750	1, 8	Q3W	6	
Vincristine	1.4	1	Q3W	6	
Prednisone	100 mg	1-5	Q3W	6	

CDOP (Cyclophosphamide, Lipo-Doxorubicin, Vincristine, Prednisone) + Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W	6-8	7, 8
Cyclophosphamide	750	1	Q3W	6-8	
Lipo-Doxorubicin	30	1	Q3W	6-8	
Vincristine	1.4	1	Q3W	6-8	
Prednisone	60	1-5	Q3W	6-8	

Concurrent presentation with CNS disease

Parenchymal

3 g/m² or more of systemic Methotrexate given on Day 15 of a 21-day RCHOP cycle that has been supported by growth factors.

Leptomeningeal

IT methotrexate/cytarabine, consider Ommaya reservoir placement and/or systemic methotrexate (3-3.5 g/m²)

Second-line Therapy and Subsequent Therapy (intention to proceed to high-dose therapy)

DHAP (Dexamethasone, Cisplatin, Cytarabine) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Cisplatin	100	1	Q3-4W		12
Cytarabine	2000 Q12H	2	Q3-4W		
Dexamethasone	40 mg	1-4	Q3-4W		

DHAX (dexamethasone, cytarabine, oxaliplatin) ± rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Oxaliplatin	100	1	Q3-4W		26
Cytarabine	2000 Q12H	2	Q3-4W		
Dexamethasone	40 mg	1-4	Q3-4W		

ESHAP (Etoposide, Methylprednisolone, Cytarabine, Cisplatin) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1	Q3-4W		13
Etoposide	40	1-4	Q3-4W		
Methylprednisolone	500 mg	1-4	Q3-4W		
Cytarabine	2000	5	Q3-4W		
Cisplatin	25	1-4	Q3-4W		

GDP (Gemcitabine, Dexamethasone, Cisplatin) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	8	Q3W		20
Gemcitabine	1000	1, 8	Q3W		
Dexamethasone	40 mg	1-4	Q3W		
Cisplatin	75	1	Q3W		

GDP (Gemcitabine, Dexamethasone, Cisplatin) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	8	Q3W		14
Gemcitabine	1000	1, 8	Q3W		
Dexamethasone	40 mg	1-4	Q3W		
Carboplatin	AUC 5	1	Q3W		

GemOx (Gemcitabine, Oxaliplatin) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1	Q2-3W		15, 23
Gemcitabine	1000	2	Q2-3W		
Oxaliplatin	100	2	Q2-3W		

ICE (Ifosfamide, Carboplatin, Etoposide) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1	Q2W		12
Etoposide	100	1-3	Q2W		
Carboplatin	AUC 5	2	Q2W		
Ifosfamide	5000	2	Q2W		

MINE (Mesna, Ifosfamide, Mitoxatrone, Etoposide) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1	Q3-4W		17
Mesna	1330	1-3	Q3-4W		
Ifosfamide	1330	1-3	Q3-4W		
Mitoxantrone	8	1	Q3-4W		
Etoposide	65	1-3	Q3-4W		

Second-line Therapy (non-candidates for high-dose therapy)

Bendamustine ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1	Q3W		18
Bendamustine	120	1-2	Q3W		

CEOP (Cyclophosphamide, Etoposide, Vincristine, Prednisone) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1	Q3W		16
Cyclophosphamide	750	1	Q3W		
Etoposide	50	1	Q3W		
Etoposide	100 PO	2-3	Q3W		
Vincristine	1.4	1	Q3W		
Prednisone	100	1-5	Q3W		

DA-EPOCH (Etoposide, Prednisone, Vincristine, Cyclophosphamide, Doxorubicin) + Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	Q3W		19
Etoposide	50	1-4	Q3W		
Doxorubicin	10	1-4	Q3W		
Vincristine	0.4	1-4	Q3W		
Cyclophosphamide	750	5	Q3W		
Prednisone	60	1-5	Q3W		

GDP (Gemcitabine, Dexamethasone, Cisplatin) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	8	Q3W		20
Gemcitabine	1000	1, 8	Q3W		
Dexamethasone	40 mg	1-4	Q3W		
Cisplatin	75	1	Q3W		

GDP (Gemcitabine, Dexamethasone, Cisplatin) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	8	Q3W		14
Gemcitabine	1000	1, 8	Q3W		
Dexamethasone	40 mg	1-4	Q3W		
Carboplatin	AUC 5	1	Q3W		

GemOx (Gemcitabine, Oxaliplatin) ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1	Q2W		21
Gemcitabine	1000-1200	1	Q2W		
Oxaliplatin	100-120	2	Q2W		

Gemcitabine, Vinorelbine ± Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1	Q3W	6	28
Gemcitabine	1000	1, 8	Q3W	6	
Vinorelbine	30	1, 8	Q3W	6	

Lenalidomide ± Rituximab (non-GCB DLBCL)

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
± Rituximab	375	1	Q4W		27
Lenalidomide	20	1-21	Q4W		

Rituximab

藥品名	劑量 mg/m ²	給藥日	頻率	週期	參考文獻
Rituximab	375	1	QW		22

★三院有個別版本

參考文獻

1. Coiffier B, Thieblemont C, Van Den Neste E, et al. Long-term outcome of patients in the LNH-98.5 trial, the first randomized study comparing rituximab-CHOP to standard CHOP chemotherapy in DLBCL patients: a study by the Groupe d'Etudes des Lymphomes de l'Adulte. *Blood* 2010;116:2040-2045.
2. Feugier P, Van hoof A, Sebban C, et al. Long-term results of the R-CHOP study in the treatment of elderly patients with diffuse large B-cell lymphoma: a study by the Groupe d'Etude des Lymphomes de l'Adulte. *J Clin Oncol* 2005;23:4117-4126.
3. Pfreundschuh M, Trumper L, Osterborg A, et al. CHOP-like chemotherapy plus rituximab versus CHOP-like chemotherapy alone in young patients with good-prognosis diffuse large-B-cell lymphoma: a randomized controlled trial by the MabThera International Trial (MInT) Group. *Lancet Oncol* 2006; 7:379-391.
4. Pfreundschuh M, Schubert J, Ziepert M, et al. Six versus eight cycles of bi-weekly CHOP-14 with or without rituximab in elderly patients with aggressive CD20+ B-cell lymphomas: a randomized controlled trial (RICOVER-60). *Lancet Oncol* 2008;9:105-116.

5. Purroy N, Bergua J, Gallur L, et al. Long-term follow-up of dose-adjusted EPOCH plus rituximab (DA-EPOCH-R) in untreated patients with poor prognosis large B-cell lymphoma. A phase II study conducted by the Spanish PETHEMA Group. *Br J Haematol* 2015;169:188-198.
6. EPOCH-rituximab in untreated diffuse large B-cell lymphoma with analysis of outcome by molecular subtype. *Haematologica* 2012;97:758-765.
7. Martino R, Perea G, Caballero MD, et al. Cyclophosphamide, pegylated liposomal doxorubicin (Caelyx), vincristine and prednisone (CCOP) in elderly patients with diffuse large B-cell lymphoma: Results from a prospective phase II study. *Haematologica* 2002;87:822-827.
8. Zaja F, Tomadini V, Zaccaria A, et al. CHOP-rituximab with pegylated liposomal doxorubicin for the treatment of elderly patients with diffuse large B-cell lymphoma. *Leuk Lymphoma* 2006; 47:2174-2180.
9. Moccia A, Schaff K, Hoskins P, et al. R-CHOP with etoposide substituted for doxorubicin (R-CEOP): Excellent outcome in diffuse large B cell lymphoma for patients with a contraindication to anthracyclines [abstract]. *Blood* 2009;114:Abstract 408.
10. Fields PA, Townsend W, Webb A, et al. De novo treatment of diffuse large B-cell lymphoma with rituximab, cyclophosphamide, vincristine, gemcitabine, and prednisolone in patients with cardiac comorbidity: a United Kingdom National Cancer Research Institute trial. *J Clin Oncol* 2014;32:282-287.
11. Peyrade F, Jardin F, Thieblemont C, et al. Attenuated immunochemotherapy regimen (R-miniCHOP) in elderly patients older than 80 years with diffuse large B-cell lymphoma: a multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2011;12:460-468.
12. Gisselbrecht C, Glass B, Mounier N, et al. Salvage regimens with autologous transplantation for relapsed large B-cell lymphoma in the rituximab era. *J Clin Oncol* 2010;28:4184-4190.
13. Martin A, Conde E, Arnan M, et al. R-ESHAP as salvage therapy for patients with relapsed or refractory diffuse large B-cell lymphoma: the influence of prior exposure to rituximab on outcome. A GEL/TAMO study. *Haematologica* 2008;93:1829-1836.
14. Gopal AK, Press OW, Shustov AR, et al. Efficacy and safety of gemcitabine, carboplatin, dexamethasone, and rituximab in patients with relapsed/refractory lymphoma: a prospective multicenter phase II study by the Puget Sound Oncology Consortium. *Leuk Lymphoma* 2010;51:1523-1529.
15. Lopez A, Gutierrez A, Palacios A, et al. GEMOX-R regimen is a highly effective salvage regimen in patients with refractory/relapsing diffuse large-cell lymphoma: a phase II study. *Eur J Haematol* 2008;80:127-132.

16. Gisselbrecht C, Glass B, Mounier N, et al. Salvage regimens with autologous transplantation for relapsed large B-cell lymphoma in the rituximab era. *J Clin Oncol* 2010;28:4184-4190.
17. Rodriguez MA, Cabanillas FC, Hagemester FB, et al. A phase II trial of mesna/ ifosfamide, mitoxantrone and etoposide for refractory lymphomas. *Ann Oncol* 1995;6(6):609-611.
18. Vacirca JL, Acs PI, Tabbara IA, et al. Bendamustine combined with rituximab for patients with relapsed or refractory diffuse large B cell lymphoma. *Ann Hematol* 2014;93:403-409.
19. Wilson WH, Jung SH, Porcu P, et al. A Cancer and Leukemia Group B multi-center study of DA-EPOCH- rituximab in untreated diffuse large B-cell lymphoma with analysis of outcome by molecular subtype. *Haematologica* 2012;97:758-765.
20. Crump M, Baetz T, Couban S, et al. Gemcitabine, dexamethasone, and cisplatin in patients with recurrent or refractory aggressive histology B-cell non-Hodgkin lymphoma: a Phase II study by the National Cancer Institute of Canada Clinical Trials Group (NCIC-CTG). *Cancer* 2004;101:1835-1842.
21. Corazzelli G, Capobianco G, Arcamone M, et al. Long-term results of gemcitabine plus oxaliplatin with and without rituximab as salvage treatment for transplant-ineligible patients with refractory/relapsing B-cell lymphoma. *Cancer Chemother Pharmacol* 2009;64:907-916.
22. Maloney DG, Grillo Lopez AJ, White CA, et al. IDEC-C2B8 (Rituximab) Anti-CD20 Monoclonal Antibody Therapy in Patients With Relapsed Low-Grade Non-Hodgkin's Lymphoma. *Blood* 1997;90(6):2188-2195.
23. El Gnaoui T, Dupuis J, Belhadj K, et al. Rituximab, gemcitabine and oxaliplatin: An effective salvage regimen for patients with relapsed or refractory B-cell lymphoma not candidates for high-dose therapy. *Ann Oncol* 2007;18:1363-1368.
24. Hiroaki A, Ipei S, Yasufumi M, et al. 3A Comparison between R-THP-COP and R-CHOP Regimens for the Treatment of Diffuse Large B-cell Lymphoma in Old Patients: A Single-institution Analysis. *Intern Med* August 21, 2017.
25. Laribi K, Denizon N, Bolle D, et al. R-CVP regimen is active in frail elderly patients aged 80 or over with diffuse large B cell lymphoma. *Ann Hematol.* 2016;95(10):1705–1714.
26. Julie Lignon, David Sibon, et al. Rituximab, Dexamethasone, Cytarabine, and Oxaliplatin (R-DHAX) Is an Effective and Safe Salvage Regimen in Relapsed/Refractory B-Cell Non-Hodgkin Lymphoma. *Clinical Lymphoma, Myeloma & Leukemia*, Vol. 10, No. 4, 262-269, 2010.

27. Wang M, Fowler N, Wagner-Bartak N, et al. Oral lenalidomide with rituximab in relapsed or refractory diffuse large cell, follicular, and transformed lymphoma: a phase II clinical trial. *Leukemia*. 2013;27:1902-1909.
28. Papageorgiou ES, Tsigotis P, Dimopoulos M, et al. Combination chemotherapy with gemcitabine and vinorelbine in the treatment of relapsed or refractory diffuse large B-cell lymphoma: a phase-II trial by the Hellenic Cooperative Oncology Group. *Eur J Hematol*. 2005;75:124-129.